

**CORYNEBACTERIUM GLUTAMICUM GENES ENCODING STRESS,
RESISTANCE AND TOLERANCE PROTEINS**

Related Applications

5 This application claims priority to prior filed U.S. Provisional Patent Application
Serial No. 60/141031, filed June 25, 1999, U.S. Provisional Patent Application Serial
No. 60/142692, filed July 1, 1999, and also to U.S. Provisional Patent Application Serial
No. 60/151214, filed August 27, 1999. This application also claims priority to German
Patent Application No. 19930429.7, filed July 1, 1999, German Patent Application No.
10 19931413.6, filed July 8, 1999, German Patent Application No. 19931457.8, filed July
8, 1999, German Patent Application No. 19931541.8, filed July 8, 1999, German Patent
Application No. 19932209.0, filed July 9, 1999, German Patent Application No.
19932230.9, filed July 9, 1999, German Patent Application No. 19932914.1, filed July
14, 1999, German Patent Application No. 19940764.9, filed August 27, 1999, and
15 German Patent Application No. 19941382.7, filed August 31, 1999. The entire contents
of all of the aforementioned applications are hereby expressly incorporated herein in
their entirety by this reference.

Background of the Invention

20 Certain products and by-products of naturally-occurring metabolic processes in
cells have utility in a wide array of industries, including the food, feed, cosmetics, and
pharmaceutical industries. These molecules, collectively termed 'fine chemicals',
include organic acids, both proteinogenic and non-proteinogenic amino acids,
nucleotides and nucleosides, lipids and fatty acids, diols, carbohydrates, aromatic
25 compounds, vitamins and cofactors, and enzymes. Their production is most
conveniently performed through large-scale culture of bacteria developed to produce
and secrete large quantities of a particular desired molecule. One particularly useful
organism for this purpose is *Corynebacterium glutamicum*, a gram positive,
nonpathogenic bacterium. Through strain selection, a number of mutant strains have
30 been developed which produce an array of desirable compounds. However, selection of
strains improved for the production of a particular molecule is a time-consuming and
difficult process.

Summary of the Invention

35 The invention provides novel bacterial nucleic acid molecules which have a
variety of uses. These uses include the identification of microorganisms which can be
used to produce fine chemicals, the modulation of fine chemical production in *C.*

glutamicum or related bacteria, the typing or identification of *C. glutamicum* or related bacteria, as reference points for mapping the *C. glutamicum* genome, and as markers for transformation. These novel nucleic acid molecules encode proteins, referred to herein as stress, resistance and tolerance (SRT) proteins.

5 *C. glutamicum* is a gram positive, aerobic bacterium which is commonly used in industry for the large-scale production of a variety of fine chemicals, and also for the degradation of hydrocarbons (such as in petroleum spills) and for the oxidation of terpenoids. The SRT nucleic acid molecules of the invention, therefore, can be used to identify microorganisms which can be used to produce fine chemicals, *e.g.*, by
10 fermentation processes. Modulation of the expression of the SRT nucleic acids of the invention, or modification of the sequence of the SRT nucleic acid molecules of the invention, can be used to modulate the production of one or more fine chemicals from a microorganism (*e.g.*, to improve the yield or production of one or more fine chemicals from a *Corynebacterium* or *Brevibacterium* species).

15 The SRT nucleic acids of the invention may also be used to identify an organism as being *Corynebacterium glutamicum* or a close relative thereof, or to identify the presence of *C. glutamicum* or a relative thereof in a mixed population of microorganisms. The invention provides the nucleic acid sequences of a number of *C. glutamicum* genes; by probing the extracted genomic DNA of a culture of a unique or
20 mixed population of microorganisms under stringent conditions with a probe spanning a region of a *C. glutamicum* gene which is unique to this organism, one can ascertain whether this organism is present. Although *Corynebacterium glutamicum* itself is nonpathogenic, it is related to species pathogenic in humans, such as *Corynebacterium diphtheriae* (the causative agent of diphtheria); the detection of such organisms is of
25 significant clinical relevance.

 The SRT nucleic acid molecules of the invention may also serve as reference points for mapping of the *C. glutamicum* genome, or of genomes of related organisms. Similarly, these molecules, or variants or portions thereof, may serve as markers for genetically engineered *Corynebacterium* or *Brevibacterium* species.

30 The SRT proteins encoded by the novel nucleic acid molecules of the invention are capable of, for example, permitting *C. glutamicum* to survive in a setting which is either chemically or environmentally hazardous to this microorganism. Given the availability of cloning vectors for use in *Corynebacterium glutamicum*, such as those disclosed in Sinskey *et al.*, U.S. Patent No. 4,649,119, and techniques for genetic
35 manipulation of *C. glutamicum* and the related *Brevibacterium* species (*e.g.*, *lactofermentum*) (Yoshihama *et al.*, *J. Bacteriol.* 162: 591-597 (1985); Katsumata *et al.*, *J. Bacteriol.* 159: 306-311 (1984); and Santamaria *et al.*, *J. Gen. Microbiol.* 130: 2237-

09603208 "062300

2246 (1984)), the nucleic acid molecules of the invention may be utilized in the genetic engineering of this organism to make it a better or more efficient producer of one or more fine chemicals, through the ability of these proteins to permit growth and multiplication of *C. glutamicum* (and also continuous production of one or more fine chemicals) under circumstances which would normally impede growth of the organism, such as those conditions frequently encountered during large-scale fermentative growth. For example, by overexpressing or engineering a heat-shock induced protease molecule such that it is optimized in activity, one may increase the ability of the bacterium to degrade incorrectly folded proteins when the bacterium is challenged with high temperatures. By having fewer misfolded (and possibly misregulated or nonfunctional) proteins to interfere with normal reaction mechanisms in the cell, the cell is increased in its ability to function normally in such a culture, which should in turn provide increased viability. This overall increase in number of cells having greater viability and activity in the culture should also result in an increase in yield, production, and/or efficiency of production of one or more desired fine chemicals, due at least to the relatively greater number of cells producing these chemicals in the culture.

This invention provides novel SRT nucleic acid molecules which encode SRT proteins which are capable of, for example, permitting *C. glutamicum* to survive in a setting which is either chemically or environmentally hazardous to this microorganism. Nucleic acid molecules encoding an SRT protein are referred to herein as SRT nucleic acid molecules. In a preferred embodiment, the SRT protein participates in metabolic pathways permitting *C. glutamicum* to survive in a setting which is either chemically or environmentally hazardous to this microorganism. Examples of such proteins include those encoded by the genes set forth in Table 1.

Accordingly, one aspect of the invention pertains to isolated nucleic acid molecules (*e.g.*, cDNAs, DNAs, or RNAs) comprising a nucleotide sequence encoding an SRT protein or biologically active portions thereof, as well as nucleic acid fragments suitable as primers or hybridization probes for the detection or amplification of SRT-encoding nucleic acid (*e.g.*, DNA or mRNA). In particularly preferred embodiments, the isolated nucleic acid molecule comprises one of the nucleotide sequences set forth in Appendix A or the coding region or a complement thereof of one of these nucleotide sequences. In other particularly preferred embodiments, the isolated nucleic acid molecule of the invention comprises a nucleotide sequence which hybridizes to or is at least about 50%, preferably at least about 60%, more preferably at least about 70%, 80% or 90%, and even more preferably at least about 95%, 96%, 97%, 98%, 99% or more homologous to a nucleotide sequence set forth in Appendix A, or a portion thereof. In other preferred embodiments, the isolated nucleic acid molecule encodes one of the

amino acid sequences set forth in Appendix B. The preferred SRT proteins of the present invention also preferably possess at least one of the SRT activities described herein.

5 In another embodiment, the isolated nucleic acid molecule encodes a protein or portion thereof wherein the protein or portion thereof includes an amino acid sequence which is sufficiently homologous to an amino acid sequence of Appendix B, *e.g.*, sufficiently homologous to an amino acid sequence of Appendix B such that the protein or portion thereof maintains an SRT activity. Preferably, the protein or portion thereof encoded by the nucleic acid molecule maintains the ability to increase the survival of *C. glutamicum* in a setting which is either chemically or environmentally hazardous to this microorganism. In one embodiment, the protein encoded by the nucleic acid molecule is at least about 50%, preferably at least about 60%, and more preferably at least about 70%, 80%, or 90% and most preferably at least about 95%, 96%, 97%, 98%, or 99% or more homologous to an amino acid sequence of Appendix B (*e.g.*, an entire amino acid sequence selected from those sequences set forth in Appendix B). In another preferred 10 embodiment, the protein is a full length *C. glutamicum* protein which is substantially homologous to an entire amino acid sequence of Appendix B (encoded by an open reading frame shown in Appendix A).

15 In another preferred embodiment, the isolated nucleic acid molecule is derived from *C. glutamicum* and encodes a protein (*e.g.*, an SRT fusion protein) which includes a biologically active domain which is at least about 50% or more homologous to one of the amino acid sequences of Appendix B and has the ability to increase the survival of *C. glutamicum* in a setting which is either chemically or environmentally hazardous to this microorganism, or possesses one or more of the activities set forth in Table 1, and 20 which also includes heterologous nucleic acid sequences encoding a heterologous polypeptide or regulatory regions.

25 In another embodiment, the isolated nucleic acid molecule is at least 15 nucleotides in length and hybridizes under stringent conditions to a nucleic acid molecule comprising a nucleotide sequence of Appendix A. Preferably, the isolated nucleic acid molecule corresponds to a naturally-occurring nucleic acid molecule. More 30 preferably, the isolated nucleic acid encodes a naturally-occurring *C. glutamicum* SRT protein, or a biologically active portion thereof.

Another aspect of the invention pertains to vectors, *e.g.*, recombinant expression vectors, containing the nucleic acid molecules of the invention, and host cells into which 35 such vectors have been introduced. In one embodiment, such a host cell is used to produce an SRT protein by culturing the host cell in a suitable medium. The SRT protein can be then isolated from the medium or the host cell.

09603208.062300

Yet another aspect of the invention pertains to a genetically altered microorganism in which an SRT gene has been introduced or altered. In one embodiment, the genome of the microorganism has been altered by the introduction of a nucleic acid molecule of the invention encoding wild-type or mutated SRT sequence as a transgene. In another embodiment, an endogenous SRT gene within the genome of the microorganism has been altered, *e.g.*, functionally disrupted, by homologous recombination with an altered SRT gene. In another embodiment, an endogenous or introduced SRT gene in a microorganism has been altered by one or more point mutations, deletions, or inversions, but still encodes a functional SRT protein. In still another embodiment, one or more of the regulatory regions (*e.g.*, a promoter, repressor, or inducer) of a SRT gene in a microorganism has been altered (*e.g.*, by deletion, truncation, inversion, or point mutation) such that the expression of the SRT gene is modulated. In a preferred embodiment, the microorganism belongs to the genus *Corynebacterium* or *Brevibacterium*, with *Corynebacterium glutamicum* being particularly preferred. In a preferred embodiment, the microorganism is also utilized for the production of a desired compound, such as an amino acid, with lysine being particularly preferred.

In another aspect, the invention provides a method of identifying the presence or activity of *Corynebacterium diphtheriae* in a subject. This method includes detection of one or more of the nucleic acid or amino acid sequences of the invention (*e.g.*, the sequences set forth in Appendix A or Appendix B) in a subject, thereby detecting the presence or activity of *Corynebacterium diphtheriae* in the subject.

Still another aspect of the invention pertains to an isolated SRT protein or a portion, *e.g.*, a biologically active portion, thereof. In a preferred embodiment, the isolated SRT protein or portion thereof possesses the ability to increase the survival of *C. glutamicum* in a setting which is either chemically or environmentally hazardous to this microorganism. In another preferred embodiment, the isolated SRT protein or portion thereof is sufficiently homologous to an amino acid sequence of Appendix B such that the protein or portion thereof maintains the ability to increase the survival of *C. glutamicum* in a setting which is either chemically or environmentally hazardous to this microorganism.

The invention also provides an isolated preparation of an SRT protein. In preferred embodiments, the SRT protein comprises an amino acid sequence of Appendix B. In another preferred embodiment, the invention pertains to an isolated full length protein which is substantially homologous to an entire amino acid sequence of Appendix B (encoded by an open reading frame set forth in Appendix A). In yet another embodiment, the protein is at least about 50%, preferably at least about 60%, and more

09603208-062300

preferably at least about 70%, 80%, or 90%, and most preferably at least about 95%, 96%, 97%, 98%, or 99% or more homologous to an entire amino acid sequence of Appendix B. In other embodiments, the isolated SRT protein comprises an amino acid sequence which is at least about 50% or more homologous to one of the amino acid sequences of Appendix B and is able to improve the survival rate of *C. glutamicum* in a setting which is either chemically or environmentally hazardous to this microorganism, or has one or more of the activities set forth in Table 1.

Alternatively, the isolated SRT protein can comprise an amino acid sequence which is encoded by a nucleotide sequence which hybridizes, *e.g.*, hybridizes under stringent conditions, or is at least about 50%, preferably at least about 60%, more preferably at least about 70%, 80%, or 90%, and even more preferably at least about 95%, 96%, 97%, 98%, or 99% or more homologous, to a nucleotide sequence of Appendix B. It is also preferred that the preferred forms of SRT proteins also have one or more of the SRT bioactivities described herein.

The SRT polypeptide, or a biologically active portion thereof, can be operatively linked to a non-SRT polypeptide to form a fusion protein. In preferred embodiments, this fusion protein has an activity which differs from that of the SRT protein alone. In other preferred embodiments, this fusion protein results in increased yields, production, and/or efficiency of production of a desired fine chemical from *C. glutamicum*. In particularly preferred embodiments, integration of this fusion protein into a host cell modulates the production of a desired compound from the cell.

In another aspect, the invention provides methods for screening molecules which modulate the activity of an SRT protein, either by interacting with the protein itself or a substrate or binding partner of the SRT protein, or by modulating the transcription or translation of an SRT nucleic acid molecule of the invention.

Another aspect of the invention pertains to a method for producing a fine chemical. This method involves the culturing of a cell containing a vector directing the expression of an SRT nucleic acid molecule of the invention, such that a fine chemical is produced. In a preferred embodiment, this method further includes the step of obtaining a cell containing such a vector, in which a cell is transfected with a vector directing the expression of an SRT nucleic acid. In another preferred embodiment, this method further includes the step of recovering the fine chemical from the culture. In a particularly preferred embodiment, the cell is from the genus *Corynebacterium* or *Brevibacterium*, or is selected from those strains set forth in Table 3.

Another aspect of the invention pertains to methods for modulating production of a molecule from a microorganism. Such methods include contacting the cell with an agent which modulates SRT protein activity or SRT nucleic acid expression such that a

00503208.062300

cell associated activity is altered relative to this same activity in the absence of the agent. In a preferred embodiment, the cell is modulated in resistance to one or more toxic chemicals or in resistance to one or more environmental stresses, such that the yields or rate of production of a desired fine chemical by this microorganism is

5 improved. The agent which modulates SRT protein activity can be an agent which stimulates SRT protein activity or SRT nucleic acid expression. Examples of agents which stimulate SRT protein activity or SRT nucleic acid expression include small molecules, active SRT proteins, and nucleic acids encoding SRT proteins that have been introduced into the cell. Examples of agents which inhibit SRT activity or expression
10 include small molecules, and antisense SRT nucleic acid molecules.

Another aspect of the invention pertains to methods for modulating yields of a desired compound from a cell, involving the introduction of a wild-type or mutant SRT gene into a cell, either maintained on a separate plasmid or integrated into the genome of the host cell. If integrated into the genome, such integration can random, or it can take
15 place by homologous recombination such that the native gene is replaced by the introduced copy, causing the production of the desired compound from the cell to be modulated. In a preferred embodiment, said yields are increased. In another preferred embodiment, said chemical is a fine chemical. In a particularly preferred embodiment, said fine chemical is an amino acid. In especially preferred embodiments, said amino
20 acid is L-lysine.

Detailed Description of the Invention

The present invention provides SRT nucleic acid and protein molecules which are involved in the survival of *C. glutamicum* upon exposure of this microorganism to chemical or environmental hazards. The molecules of the invention may be utilized in
25 the modulation of production of fine chemicals from microorganisms, since these SRT proteins provide a means for continued growth and multiplication of *C. glutamicum* in the presence of toxic chemicals or hazardous environmental conditions, such as may be encountered during large-scale fermentative growth. By increasing the growth rate or at least maintaining normal growth in the face of poor, if not toxic, conditions, one may
30 increase the yield, production, and/or efficiency of production of one or more fine chemicals from such a culture, at least due to the relatively greater number of cells producing the fine chemical in the culture. Aspects of the invention are further explicated below.

I. Fine Chemicals

The term 'fine chemical' is art-recognized and includes molecules produced by an organism which have applications in various industries, such as, but not limited to, the pharmaceutical, agriculture, and cosmetics industries. Such compounds include organic acids, such as tartaric acid, itaconic acid, and diaminopimelic acid, both proteinogenic and non-proteinogenic amino acids, purine and pyrimidine bases, nucleosides, and nucleotides (as described *e.g.* in Kuninaka, A. (1996) Nucleotides and related compounds, p. 561-612, in Biotechnology vol. 6, Rehm *et al.*, eds. VCH: Weinheim, and references contained therein), lipids, both saturated and unsaturated fatty acids (*e.g.*, arachidonic acid), diols (*e.g.*, propane diol, and butane diol), carbohydrates (*e.g.*, hyaluronic acid and trehalose), aromatic compounds (*e.g.*, aromatic amines, vanillin, and indigo), vitamins and cofactors (as described in Ullmann's Encyclopedia of Industrial Chemistry, vol. A27, "Vitamins", p. 443-613 (1996) VCH: Weinheim and references therein; and Ong, A.S., Niki, E. & Packer, L. (1995) "Nutrition, Lipids, Health, and Disease" Proceedings of the UNESCO/Confederation of Scientific and Technological Associations in Malaysia, and the Society for Free Radical Research – Asia, held Sept. 1-3, 1994 at Penang, Malaysia, AOCS Press, (1995)), enzymes, polyketides (Cane *et al.* (1998) *Science* 282: 63-68), and all other chemicals described in Gutcho (1983) Chemicals by Fermentation, Noyes Data Corporation, ISBN: 0818805086 and references therein. The metabolism and uses of certain of these fine chemicals are further explicated below.

A. *Amino Acid Metabolism and Uses*

Amino acids comprise the basic structural units of all proteins, and as such are essential for normal cellular functioning in all organisms. The term "amino acid" is art-recognized. The proteinogenic amino acids, of which there are 20 species, serve as structural units for proteins, in which they are linked by peptide bonds, while the nonproteinogenic amino acids (hundreds of which are known) are not normally found in proteins (see Ullmann's Encyclopedia of Industrial Chemistry, vol. A2, p. 57-97 VCH: Weinheim (1985)). Amino acids may be in the D- or L- optical configuration, though L-amino acids are generally the only type found in naturally-occurring proteins. Biosynthetic and degradative pathways of each of the 20 proteinogenic amino acids have been well characterized in both prokaryotic and eukaryotic cells (see, for example, Stryer, L. Biochemistry, 3rd edition, pages 578-590 (1988)). The 'essential' amino acids (histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan, and valine), so named because they are generally a nutritional requirement due to the complexity of their biosyntheses, are readily converted by simple biosynthetic pathways

006290" 8020960

to the remaining 11 'nonessential' amino acids (alanine, arginine, asparagine, aspartate, cysteine, glutamate, glutamine, glycine, proline, serine, and tyrosine). Higher animals do retain the ability to synthesize some of these amino acids, but the essential amino acids must be supplied from the diet in order for normal protein synthesis to occur.

5 Aside from their function in protein biosynthesis, these amino acids are interesting chemicals in their own right, and many have been found to have various applications in the food, feed, chemical, cosmetics, agriculture, and pharmaceutical industries. Lysine is an important amino acid in the nutrition not only of humans, but also of monogastric animals such as poultry and swine. Glutamate is most commonly
10 used as a flavor additive (mono-sodium glutamate, MSG) and is widely used throughout the food industry, as are aspartate, phenylalanine, glycine, and cysteine. Glycine, L-methionine and tryptophan are all utilized in the pharmaceutical industry. Glutamine, valine, leucine, isoleucine, histidine, arginine, proline, serine and alanine are of use in both the pharmaceutical and cosmetics industries. Threonine, tryptophan, and D/ L-
15 methionine are common feed additives. (Leuchtenberger, W. (1996) Amino acids – technical production and use, p. 466-502 in Rehm *et al.* (eds.) Biotechnology vol. 6, chapter 14a, VCH: Weinheim). Additionally, these amino acids have been found to be useful as precursors for the synthesis of synthetic amino acids and proteins, such as N-acetylcysteine, S-carboxymethyl-L-cysteine, (S)-5-hydroxytryptophan, and others
20 described in Ulmann's Encyclopedia of Industrial Chemistry, vol. A2, p. 57-97, VCH: Weinheim, 1985.

 The biosynthesis of these natural amino acids in organisms capable of producing them, such as bacteria, has been well characterized (for review of bacterial amino acid biosynthesis and regulation thereof, see Umbarger, H.E.(1978) *Ann. Rev.*
25 *Biochem.* 47: 533-606). Glutamate is synthesized by the reductive amination of α -ketoglutarate, an intermediate in the citric acid cycle. Glutamine, proline, and arginine are each subsequently produced from glutamate. The biosynthesis of serine is a three-step process beginning with 3-phosphoglycerate (an intermediate in glycolysis), and resulting in this amino acid after oxidation, transamination, and hydrolysis steps. Both
30 cysteine and glycine are produced from serine; the former by the condensation of homocysteine with serine, and the latter by the transferal of the side-chain β -carbon atom to tetrahydrofolate, in a reaction catalyzed by serine transhydroxymethylase. Phenylalanine, and tyrosine are synthesized from the glycolytic and pentose phosphate pathway precursors erythrose 4-phosphate and phosphoenolpyruvate in a 9-step
35 biosynthetic pathway that differ only at the final two steps after synthesis of prephenate. Tryptophan is also produced from these two initial molecules, but its synthesis is an 11-step pathway. Tyrosine may also be synthesized from phenylalanine, in a reaction

00602208 062300

5 formed from threonine. A complex 9-step pathway results in the production of histidine from 5-phosphoribosyl-1-pyrophosphate, an activated sugar.

and the Urea Cycle” p. 495-516 (1988)). Although the cell is able to convert unwanted amino acids into useful metabolic intermediates, amino acid production is costly in terms of energy, precursor molecules, and the enzymes necessary to synthesize them. Thus it is not surprising that amino acid biosynthesis is regulated by feedback inhibition, in which the presence of a particular amino acid serves to slow or entirely stop its own production (for overview of feedback mechanisms in amino acid biosynthetic pathways, see Stryer, L. Biochemistry, 3rd ed. Ch. 24: “Biosynthesis of Amino Acids and Heme” p. 575-600 (1988)). Thus, the output of any particular amino acid is limited by the amount of that amino acid present in the cell.

20 *B. Vitamin, Cofactor, and Nutraceutical Metabolism and Uses*

substances which may serve as electron carriers or intermediates in a variety of metabolic pathways. Aside from their nutritive value, these compounds also have significant industrial value as coloring agents, antioxidants, and catalysts or other processing aids. (For an overview of the structure, activity, and industrial applications of these compounds, see, for example, Ullman's Encyclopedia of Industrial Chemistry, "Vitamins" vol. A27, p. 443-613, VCH: Weinheim, 1996.) The term "vitamin" is art-recognized, and includes nutrients which are required by an organism for normal functioning, but which that organism cannot synthesize by itself. The group of vitamins may encompass cofactors and nutraceutical compounds. The language "cofactor" includes nonproteinaceous compounds required for a normal enzymatic activity to occur. Such compounds may be organic or inorganic; the cofactor molecules of the invention are preferably organic. The term "nutraceutical" includes dietary supplements having health benefits in plants and animals, particularly humans. Examples of such

molecules are vitamins, antioxidants, and also certain lipids (*e.g.*, polyunsaturated fatty acids).

The biosynthesis of these molecules in organisms capable of producing them, such as bacteria, has been largely characterized (Ullman's Encyclopedia of Industrial Chemistry, "Vitamins" vol. A27, p. 443-613, VCH: Weinheim, 1996; Michal, G. (1999) Biochemical Pathways: An Atlas of Biochemistry and Molecular Biology, John Wiley & Sons; Ong, A.S., Niki, E. & Packer, L. (1995) "Nutrition, Lipids, Health, and Disease" Proceedings of the UNESCO/Confederation of Scientific and Technological Associations in Malaysia, and the Society for Free Radical Research – Asia, held Sept. 1-3, 1994 at Penang, Malaysia, AOCS Press: Champaign, IL X, 374 S).

Thiamin (vitamin B₁) is produced by the chemical coupling of pyrimidine and thiazole moieties. Riboflavin (vitamin B₂) is synthesized from guanosine-5'-triphosphate (GTP) and ribose-5'-phosphate. Riboflavin, in turn, is utilized for the synthesis of flavin mononucleotide (FMN) and flavin adenine dinucleotide (FAD). The family of compounds collectively termed 'vitamin B₆' (*e.g.*, pyridoxine, pyridoxamine, pyridoxa-5'-phosphate, and the commercially used pyridoxin hydrochloride) are all derivatives of the common structural unit, 5-hydroxy-6-methylpyridine. Pantothenate (pantothenic acid, (R)-(+)-N-(2,4-dihydroxy-3,3-dimethyl-1-oxobutyl)-β-alanine) can be produced either by chemical synthesis or by fermentation. The final steps in pantothenate biosynthesis consist of the ATP-driven condensation of β-alanine and pantoic acid. The enzymes responsible for the biosynthesis steps for the conversion to pantoic acid, to β-alanine and for the condensation to pantothenic acid are known. The metabolically active form of pantothenate is Coenzyme A, for which the biosynthesis proceeds in 5 enzymatic steps. Pantothenate, pyridoxal-5'-phosphate, cysteine and ATP are the precursors of Coenzyme A. These enzymes not only catalyze the formation of pantothenate, but also the production of (R)-pantoic acid, (R)-pantolacton, (R)-panthenol (provitamin B₅), pantetheine (and its derivatives) and coenzyme A.

Biotin biosynthesis from the precursor molecule pimeloyl-CoA in microorganisms has been studied in detail and several of the genes involved have been identified. Many of the corresponding proteins have been found to also be involved in Fe-cluster synthesis and are members of the nifS class of proteins. Lipoic acid is derived from octanoic acid, and serves as a coenzyme in energy metabolism, where it becomes part of the pyruvate dehydrogenase complex and the α-ketoglutarate dehydrogenase complex. The folates are a group of substances which are all derivatives of folic acid, which in turn is derived from L-glutamic acid, p-amino-benzoic acid and 6-methylpterin. The biosynthesis of folic acid and its derivatives, starting from the

Corrinoids (such as the cobalamines and particularly vitamin B₁₂) and porphyrines belong to a group of chemicals characterized by a tetrapyrrole ring system.

10 phosphate) and their reduced forms.

15 its synthesis. *In vitro* methodologies require significant inputs of materials and time,
often at great cost.

C. Purine, Pyrimidine, Nucleoside and Nucleotide Metabolism and Uses

20 important targets for the therapy of tumor diseases and viral infections. The language

25 D-deoxyribose), and phosphoric acid. The language “nucleoside” includes molecules

35 R.I. and Lyons, S.D. (1990) "Potent inhibitors of *de novo* pyrimidine and purine

biosynthesis as chemotherapeutic agents.” *Med. Res. Reviews* 10: 505-548). Studies of enzymes involved in purine and pyrimidine metabolism have been focused on the

5
10

15
20
25
30

D. Trehalose Metabolism and Uses

35

Patent No. 5,759,610; Singer, M.A. and Lindquist, S. (1998) *Trends Biotech.* 16: 460-467; Paiva, C.L.A. and Panek, A.D. (1996) *Biotech. Ann. Rev.* 2: 293-314; and Shiosaka, M. (1997) *J. Japan* 172: 97-102). Trehalose is produced by enzymes from many microorganisms and is naturally released into the surrounding medium, from which it can be collected using methods known in the art.

II. Resistance to Damage from Chemicals, Environmental Stress, and Antibiotics

Production of fine chemicals is typically performed by large-scale culture of bacteria developed to produce and secrete large quantities of these molecules. However, this type of large-scale fermentation results in the subjection of the microorganisms to stresses of various kinds. These stresses include environmental stress and chemical stress.

A. Resistance to Environmental Stress

Examples of environmental stresses typically encountered in large-scale fermentative culture include mechanical stress, heat stress, stress due to limited oxygen, stress due to oxygen radicals, pH stress, and osmotic stress. The stirring mechanism used in most large-scale fermentors to ensure aeration of the culture produces heat, thus increasing the temperature of the culture. Increases in temperature induce the well-characterized heat shock response, in which a set of proteins are expressed which not only aid in the survival of the bacterium in the face of high temperatures, but also increase survival in response to a number of other environmental stresses (see Neidhardt, F.C., *et al.*, eds. (1996) *E. coli* and *Salmonella*. ASM Press: Washington, D.C., p. 1382-1399; Wosten, M. M. (1998) *FEMS Microbiology Reviews* 22(3): 127-50; Bahl, H. *et al.* (1995) *FEMS Microbiology Reviews* 17(3): 341-348; Zimmerman, J.L., Cohill, P.R. (1991) *New Biologist* 3(7): 641-650; Samali, A., and Orrenius, S. (1998) *Cell. Stress Chaperones* 3(4): 228-236, and references contained therein from each of these citations). Regulation of the heat shock response in bacteria is facilitated by specific sigma factors and other cellular regulators of gene expression (Hecker, M., Volker, U (1998). *Molecular Microbiology* 29(5): 1129-1136). One of the largest problems that the cell encounters when exposed to high temperature is that protein folding is impaired; nascent proteins have sufficient kinetic energy in high temperature circumstances that it is difficult for the growing polypeptide chain to remain in a stable conformation long enough to fold properly. Thus, two of the key types of proteins expressed during the heat shock response consist of chaperones (proteins which assist in the folding or unfolding of other proteins – see, *e.g.*, Fink, A.L. (1999) *Physiol. Rev.* 79(2): 425-449), and proteases, which can destroy any improperly folded proteins.

Examples of chaperones expressed during the heat shock response include GroEL and DNAK; proteases known to be expressed during this cellular reaction to heat shock include Lon, FtsH, and ClpB.

Other environmental stresses besides heat may also provoke a stress response.

5 Though the fermentor stirring process is meant to introduce oxygen into the culture, oxygen may remain in limited supply, particularly when the culture is advanced in growth and the oxygen needs of the culture are thereby increased; an insufficient supply of oxygen is another stress for the microorganism. Cells in fermentor cultures are also subjected to a number of osmotic stresses, particularly when nutrients are added to the
10 culture, resulting in a high extracellular and low intracellular concentration of these molecules. Further, the large quantities of the desired molecules produced by these organisms in culture may contribute to osmotic stress of the bacteria. Lastly, aerobic metabolism such as that used by *C. glutamicum* results in carbon dioxide as a waste product; secretion of this molecule may acidify the culture medium due to conversion of
15 this molecule to carboxylic acid. Thus, bacteria in culture are also frequently subjected to acidic pH stress. The converse may also be true – when high levels of basic waste molecules such as ammonium are present in the culture medium, the bacteria in culture may be subjected to basic pH stress as well.

To combat such environmental stresses, bacteria have elegant gene systems
20 which are expressed upon exposure to one or more stresses, such as the aforementioned heat shock system. Genes expressed in response to osmotic stress, for example, encode proteins capable of transporting or synthesizing compatible solutes such that osmotic intake or export of a particular molecule is slowed to manageable levels. Other examples of stress-induced bacterial proteins are those involved in trehalose biosynthesis, those
25 encoding enzymes involved in ppGpp metabolism, those involved in signal transduction, particularly those encoding two-component systems which are sensitive to osmotic pressure, and those encoding transcription factors which are responsive to a variety of stress factors (*e.g.*, RssB analogues and/or sigma factors). Many other such genes and their protein products are known in the art.

30

B. Resistance to Chemical Stress

Aside from environmental stresses, cells may also experience a number of chemical stresses. These may fall into two categories. The first are natural waste products of metabolism and other cellular processes which are secreted by the cell to the
35 surrounding medium. The second are chemicals present in the extracellular medium which do not originate from the cell. Generally, when cells excrete toxic waste products from the concentrated intracellular cytoplasm into the relatively much more dilute

09603208-062300

extracellular medium, these products dissipate such that extracellular levels of the possibly toxic compound are quite low. However, in large-scale fermentative culture of the bacterium, this may not be the case: so many bacteria are grown in a relatively small environment and at such a high metabolic rate that waste products may accumulate in the medium to nearly toxic levels. Examples of such wastes are carbon dioxide, metal ions, and reactive oxygen species such as hydrogen peroxide. These compounds may interfere with the activity or structure of cell surface molecules, or may re-enter the cell, where they can seriously damage proteins and nucleic acids alike. Certain other chemicals hazardous to the normal functioning of cells may be naturally found in the extracellular medium. For example, metal ions such as mercury, cadmium, nickel or copper are frequently found in water sources, and may form tight complexes with cellular enzymes which prevent the normal functioning of these proteins.

C. Resistance to Antibiotics

Bacteriocidal proteins or antibiotics, may also be found in the extracellular milieu, either through the intervention of the researcher, or as a natural product from another organism, utilized to gain a competitive advantage. Microorganisms have several art-known mechanisms to protect themselves against antimicrobial chemicals. Degradation, modification, and export of compounds toxic to the cell are common methods by which microorganisms eliminate or detoxify antibiotics. Cytoplasmic 'efflux-pumps' are known in several prokaryotes and show similarities to the so-called 'multidrug resistance' proteins from higher eukaryotes (Neyfakh, A. A. , *et al.* (1991) *Proc. Natl. Acad. Sci. USA* 88: 4781-4785). Examples of such proteins include emrAB from *E. coli* (Lomovskaya, O. and K. Lewis (1992) *Proc. Natl. Acad. Sci. USA* 89: 8938-8942), lmrB from *B. subtilis* (Kumano, M. *et al.* (1997) *Microbiology* 143: 2775-2782), smr from *S. aureus* (Grinius, L.G. *et al.* (1992) *Plasmid* 27: 119-129) or cmr from *C. glutamicum* (Kaidoh, K. *et al.* (1997) *Micro. Drug Resist.* 3: 345-350). *C. glutamicum* itself is non-pathogenic, in contrast to several other members of the genus *Corynebacterium* , such as *C. diphtheriae* or *C. pseudotuberculosis*. Several pathogenic *Corynebacteria* are known to have multiple resistances against a variety of antibiotics, such as *C. jeikeium* and *C. urealyticum* (Soriano, F. *et al.* (1995) *Antimicrob. Agents Chemother.* 39: 208-214).

Lincosamides are recognized as effective antibiotics against *Corynebacterium* species (Soriano, F. *et al.* (1995) *Antimicrob. Agents Chemother.* 39: 208-214). An unexpected result of the present invention was the identification of a gene encoding a lincosamide-resistance protein (in particular, a lincomycin-resistance protein). The LMRB protein from *C. glutamicum* shows 40% homology to the product of the lmrB

10

20

25

35

The language, "SRT protein" or "SRT polypeptide" includes proteins which participate in the resistance of *C. glutamicum* to one or more environmental or chemical stresses. Examples of SRT proteins include those encoded by the SRT genes set forth in Table 1 and Appendix A. The terms "SRT gene" or "SRT nucleic acid sequence" include nucleic acid sequences encoding an SRT protein, which consist of a coding region and also corresponding untranslated 5' and 3' sequence regions. Examples of SRT genes include those set forth in Table 1. The terms "production" or "productivity" are art-recognized and include the concentration of the fermentation product (for example, the desired fine chemical) formed within a given time and a given fermentation volume (e.g., kg product per hour per liter). The term "efficiency of production" includes the time required for a particular level of production to be achieved (for example, how long it takes for the cell to attain a particular rate of output of a fine chemical). The term "yield" or "product/carbon yield" is art-recognized and includes the efficiency of the conversion of the carbon source into the product (i.e., fine chemical). This is generally written as, for example, kg product per kg carbon source. By increasing the yield or production of the compound, the quantity of recovered molecules, or of useful recovered molecules of that compound in a given amount of culture over a given amount of time is increased. The terms "biosynthesis" or a "biosynthetic pathway" are art-recognized and include the synthesis of a compound, preferably an organic compound, by a cell from intermediate compounds in what may be a multistep and highly regulated process. The terms "degradation" or a "degradation pathway" are art-recognized and include the breakdown of a compound, preferably an organic compound, by a cell to degradation products (generally speaking, smaller or less complex molecules) in what may be a multistep and highly regulated process. The language "metabolism" is art-recognized and includes the totality of the biochemical reactions that take place in an organism. The metabolism of a particular compound, then, (e.g., the metabolism of an amino acid such as glycine) comprises the overall biosynthetic, modification, and degradation pathways in the cell related to this compound. The terms "resistance" and "tolerance" are art-known and include the ability of a cell to not be affected by exposure to a chemical or an environment which would otherwise be detrimental to the normal functioning of these organisms. The terms "stress" or "hazard" include factors which are detrimental to the normal functioning of cells such as *C. glutamicum*. Examples of stresses include "chemical stress", in which a cell is exposed to one or more chemicals which are detrimental to the cell, and "environmental stress" where a cell is exposed to an environmental condition outside of those to which it is adapted. Chemical stresses may be either natural metabolic waste products such as, but not limited to reactive oxygen species or carbon dioxide, or

09603208-062300

chemicals otherwise present in the environment, including, but not limited to heavy metal ions or bacteriocidal proteins such as antibiotics. Environmental stresses may be, but are not limited to temperatures outside of the normal range, suboptimal oxygen availability, osmotic pressures, or extremes of pH, for example.

5 In another embodiment, the SRT molecules of the invention are capable of modulating the production of a desired molecule, such as a fine chemical, in a microorganism such as *C. glutamicum*. Using recombinant genetic techniques, one or more of the SRT proteins of the invention may be manipulated such that its function is modulated. The alteration of activity of stress response, resistance or tolerance genes
10 such that the cell is increased in tolerance to one or more stresses may improve the ability of that cell to grow and multiply in the relatively stressful conditions of large-scale fermentor culture. For example, by overexpressing or engineering a heat-shock induced chaperone molecule such that it is optimized in activity, one may increase the ability of the bacterium to correctly fold proteins in the face of nonoptimal temperature
15 conditions. By having fewer misfolded (and possibly misregulated or nonfunctional) proteins, the cell is increased in its ability to function normally in such a culture, which should in turn provide increased viability. This overall increase in number of cells having greater viability and activity in the culture should also result in an increase in the yield, production, and/or efficiency of production of one or more desired fine chemicals,
20 due at least to the relatively greater number of cells producing these chemicals in the culture.

 The isolated nucleic acid sequences of the invention are contained within the genome of a *Corynebacterium glutamicum* strain available through the American Type Culture Collection, given designation ATCC 13032. The nucleotide sequence of the
25 isolated *C. glutamicum* SRT DNAs and the predicted amino acid sequences of the *C. glutamicum* SRT proteins are shown in Appendices A and B, respectively. Computational analyses were performed which classified and/or identified these nucleotide sequences as sequences which encode chemical and environmental stress, resistance, and tolerance proteins.

30 The present invention also pertains to proteins which have an amino acid sequence which is substantially homologous to an amino acid sequence of Appendix B. As used herein, a protein which has an amino acid sequence which is substantially homologous to a selected amino acid sequence is least about 50% homologous to the selected amino acid sequence, *e.g.*, the entire selected amino acid sequence. A protein
35 which has an amino acid sequence which is substantially homologous to a selected amino acid sequence can also be least about 50-60%, preferably at least about 60-70%, and more preferably at least about 70-80%, 80-90%, or 90-95%, and most preferably at

00603208-062300

least about 96%, 97%, 98%, 99% or more homologous to the selected amino acid sequence. Ranges and identity values intermediate to the above-recited values, (e.g., 75%-80% identical, 85-87% identical, 91-92% identical) are also intended to be encompassed by the present invention. For example, ranges of identity values using a combination of any of the above values recited as upper and/or lower limits are intended to be included.

The SRT proteins or biologically active portions or fragments thereof of the invention can confer resistance or tolerance to one or more chemical or environmental stresses, or may have one or more of the activities set forth in Table 1.

Various aspects of the invention are described in further detail in the following subsections:

A. Isolated Nucleic Acid Molecules

One aspect of the invention pertains to isolated nucleic acid molecules that encode SRT polypeptides or biologically active portions thereof, as well as nucleic acid fragments sufficient for use as hybridization probes or primers for the identification or amplification of SRT-encoding nucleic acid (e.g., SRT DNA). As used herein, the term "nucleic acid molecule" is intended to include DNA molecules (e.g., cDNA or genomic DNA) and RNA molecules (e.g., mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. This term also encompasses untranslated sequence located at both the 3' and 5' ends of the coding region of the gene: at least about 100 nucleotides of sequence upstream from the 5' end of the coding region and at least about 20 nucleotides of sequence downstream from the 3' end of the coding region of the gene. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA. An "isolated" nucleic acid molecule is one which is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated SRT nucleic acid molecule can contain less than about 5 kb, 4kb, 3kb, 2kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived (e.g., a *C. glutamicum* cell). Moreover, an "isolated" nucleic acid molecule, such as a DNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or chemical precursors or other chemicals when chemically synthesized.

In a preferred embodiment, an isolated nucleic acid molecule of the invention comprises one of the nucleotide sequences shown in Appendix A. The sequences of Appendix A correspond to the *Corynebacterium glutamicum* SRT DNAs of the invention. This DNA comprises sequences encoding SRT proteins (*i.e.*, the "coding region", indicated in each sequence in Appendix A), as well as 5' untranslated sequences and 3' untranslated sequences, also indicated in Appendix A. Alternatively, the nucleic acid molecule can comprise only the coding region of any of the sequences in Appendix

35 A. For the purposes of this application, it will be understood that each of the sequences set forth in Appendix A has an identifying RXA, RXN, or RXS number having the designation "RXA", "RXN", or "RXS" followed by 5 digits (*i.e.*,

5
10
15

20

25

30

35

09603208-062300
006290-8026960

In still another preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleotide sequence which is at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, or 60%, preferably at least about 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, or 70%, more preferably at least about 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, or 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, or 90%, or 91%, 92%, 93%, 94%, and even more preferably at least about 95%, 96%, 97%, 98%, 99% or more homologous to a nucleotide sequence shown in Appendix A, or a portion thereof. Ranges and identity values intermediate to the above-recited ranges, (e.g., 70-90% identical or 80-95% identical) are also intended to be encompassed by the present invention. For example, ranges of identity values using a combination of any of the above values recited as upper and/or lower limits are intended to be included. In an additional preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleotide sequence which hybridizes, e.g., hybridizes under stringent conditions, to one of the nucleotide sequences shown in Appendix A, or a portion thereof.

Moreover, the nucleic acid molecule of the invention can comprise only a portion of the coding region of one of the sequences in Appendix A, for example a fragment which can be used as a probe or primer or a fragment encoding a biologically active portion of an SRT protein. The nucleotide sequences determined from the cloning of the SRT genes from *C. glutamicum* allows for the generation of probes and primers designed for use in identifying and/or cloning SRT homologues in other cell types and organisms, as well as SRT homologues from other *Corynebacteria* or related species. The probe/primer typically comprises substantially purified oligonucleotide. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, preferably about 25, more preferably about 40, 50 or 75 consecutive nucleotides of a sense strand of one of the sequences set forth in Appendix A, an anti-sense sequence of one of the sequences set forth in Appendix A, or naturally occurring mutants thereof. Primers based on a nucleotide sequence of Appendix A can be used in PCR reactions to clone SRT homologues. Probes based on the SRT nucleotide sequences can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. In preferred embodiments, the probe further comprises a label group attached thereto, e.g. the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells which misexpress an SRT protein, such as by measuring a level of an SRT-encoding nucleic acid in a sample of cells, e.g., detecting SRT mRNA levels or determining whether a genomic SRT gene has been mutated or deleted.

00603208.062300

006290" 80260960

In one embodiment, the nucleic acid molecule of the invention encodes a protein or portion thereof which includes an amino acid sequence which is sufficiently homologous to an amino acid sequence of Appendix B such that the protein or portion thereof maintains the ability to confer resistance or tolerance of *C. glutamicum* to one or more chemical or environmental stresses. As used herein, the language "sufficiently homologous" refers to proteins or portions thereof which have amino acid sequences which include a minimum number of identical or equivalent (*e.g.*, an amino acid residue which has a similar side chain as an amino acid residue in one of the sequences of Appendix B) amino acid residues to an amino acid sequence of Appendix B such that the protein or portion thereof is capable of participating in the resistance of *C. glutamicum* to one or more chemical or environmental stresses. Protein members of such metabolic pathways, as described herein, function to increase the resistance or tolerance of *C. glutamicum* to one or more environmental or chemical hazards or stresses. Examples of such activities are also described herein. Thus, "the function of an SRT protein" contributes to the overall resistance of *C. glutamicum* to elements of its surroundings which may impede its normal growth or functioning, and/or contributes, either directly or indirectly, to the yield, production, and/or efficiency of production of one or more fine chemicals. Examples of SRT protein activities are set forth in Table 1.

In another embodiment, the protein is at least about 50-60%, preferably at least about 60-70%, and more preferably at least about 70-80%, 80-90%, 90-95%, and most preferably at least about 96%, 97%, 98%, 99% or more homologous to an entire amino acid sequence of Appendix B. Ranges and identity values intermediate to the above-recited values, (*e.g.*, 75%-80% identical, 85-87% identical, or 91-92% identical) are also intended to be encompassed by the present invention. For example, ranges of identity values using a combination of any of the above values recited as upper and/or lower limits are intended to be included.

Portions of proteins encoded by the SRT nucleic acid molecules of the invention are preferably biologically active portions of one of the SRT proteins. As used herein, the term "biologically active portion of an SRT protein" is intended to include a portion, *e.g.*, a domain/motif, of an SRT protein that is capable of imparting resistance or tolerance to one or more environmental or chemical stresses or hazards, or has an activity as set forth in Table 1. To determine whether an SRT protein or a biologically active portion thereof can increase the resistance or tolerance of *C. glutamicum* to one or more chemical or environmental stresses or hazards, an assay of enzymatic activity may be performed. Such assay methods are well known to those of ordinary skill in the art, as detailed in Example 8 of the Exemplification.

Additional nucleic acid fragments encoding biologically active portions of an SRT protein can be prepared by isolating a portion of one of the sequences in Appendix B, expressing the encoded portion of the SRT protein or peptide (*e.g.*, by recombinant expression *in vitro*) and assessing the activity of the encoded portion of the SRT protein or peptide.

The invention further encompasses nucleic acid molecules that differ from one of the nucleotide sequences shown in Appendix A (and portions thereof) due to degeneracy of the genetic code and thus encode the same SRT protein as that encoded by the nucleotide sequences shown in Appendix A. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence shown in Appendix B. In a still further embodiment, the nucleic acid molecule of the invention encodes a full length *C. glutamicum* protein which is substantially homologous to an amino acid sequence of Appendix B (encoded by an open reading frame shown in Appendix A).

It will be understood by one of ordinary skill in the art that in one embodiment the sequences of the invention are not meant to include the sequences of the prior art, such as those Genbank sequences set forth in Tables 2 or 4 which were available prior to the present invention. In one embodiment, the invention includes nucleotide and amino acid sequences having a percent identity to a nucleotide or amino acid sequence of the invention which is greater than that of a sequence of the prior art (*e.g.*, a Genbank sequence (or the protein encoded by such a sequence) set forth in Tables 2 or 4). For example, the invention includes a nucleotide sequence which is greater than and/or at least 39% identical to the nucleotide sequence designated RXA00084 (SEQ ID NO:189), a nucleotide sequence which is greater than and/or at least 56% identical to the nucleotide sequence designated RXA00605 (SEQ ID NO:11), and a nucleotide sequence which is greater than and/or at least 50% identical to the nucleotide sequence designated RXA00886 (SEQ ID NO:39). One of ordinary skill in the art would be able to calculate the lower threshold of percent identity for any given sequence of the invention by examining the GAP-calculated percent identity scores set forth in Table 4 for each of the three top hits for the given sequence, and by subtracting the highest GAP-calculated percent identity from 100 percent. One of ordinary skill in the art will also appreciate that nucleic acid and amino acid sequences having percent identities greater than the lower threshold so calculated (*e.g.*, at least 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, or 60%, preferably at least about 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, or 70%, more preferably at least about 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, or 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, or

90%, or 91%, 92%, 93%, 94%, and even more preferably at least about 95%, 96%, 97%, 98%, 99% or more identical) are also encompassed by the invention.

In addition to the *C. glutamicum* SRT nucleotide sequences shown in Appendix A, it will be appreciated by one of ordinary skill in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequences of SRT proteins may exist within a population (e.g., the *C. glutamicum* population). Such genetic polymorphism in the SRT gene may exist among individuals within a population due to natural variation. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame encoding an SRT protein, preferably a *C. glutamicum* SRT protein. Such natural variations can typically result in 1-5% variance in the nucleotide sequence of the SRT gene. Any and all such nucleotide variations and resulting amino acid polymorphisms in SRT that are the result of natural variation and that do not alter the functional activity of SRT proteins are intended to be within the scope of the invention.

Nucleic acid molecules corresponding to natural variants and non-*C. glutamicum* homologues of the *C. glutamicum* SRT DNA of the invention can be isolated based on their homology to the *C. glutamicum* SRT nucleic acid disclosed herein using the *C. glutamicum* DNA, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions. Accordingly, in another embodiment, an isolated nucleic acid molecule of the invention is at least 15 nucleotides in length and hybridizes under stringent conditions to the nucleic acid molecule comprising a nucleotide sequence of Appendix A. In other embodiments, the nucleic acid is at least 30, 50, 100, 250 or more nucleotides in length. As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% homologous to each other typically remain hybridized to each other. Preferably, the conditions are such that sequences at least about 65%, more preferably at least about 70%, and even more preferably at least about 75% or more homologous to each other typically remain hybridized to each other. Such stringent conditions are known to those of ordinary skill in the art and can be found in Ausubel *et al.*, *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. A preferred, non-limiting example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65°C. Preferably, an isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to a sequence of Appendix A corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having

a nucleotide sequence that occurs in nature (*e.g.*, encodes a natural protein). In one embodiment, the nucleic acid encodes a natural *C. glutamicum* SRT protein.

In addition to naturally-occurring variants of the SRT sequence that may exist in the population, one of ordinary skill in the art will further appreciate that changes can be introduced by mutation into a nucleotide sequence of Appendix A, thereby leading to changes in the amino acid sequence of the encoded SRT protein, without altering the functional ability of the SRT protein. For example, nucleotide substitutions leading to amino acid substitutions at "non-essential" amino acid residues can be made in a sequence of Appendix A. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence of one of the SRT proteins (Appendix B) without altering the activity of said SRT protein, whereas an "essential" amino acid residue is required for SRT protein activity. Other amino acid residues, however, (*e.g.*, those that are not conserved or only semi-conserved in the domain having SRT activity) may not be essential for activity and thus are likely to be amenable to alteration without altering SRT activity.

Accordingly, another aspect of the invention pertains to nucleic acid molecules encoding SRT proteins that contain changes in amino acid residues that are not essential for SRT activity. Such SRT proteins differ in amino acid sequence from a sequence contained in Appendix B yet retain at least one of the SRT activities described herein. In one embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence encoding a protein, wherein the protein comprises an amino acid sequence at least about 50% homologous to an amino acid sequence of Appendix B and is capable of increasing the resistance or tolerance of *C. glutamicum* to one or more environmental or chemical stresses, or has one or more of the activities set forth in Table 1. Preferably, the protein encoded by the nucleic acid molecule is at least about 50-60% homologous to one of the sequences in Appendix B, more preferably at least about 60-70% homologous to one of the sequences in Appendix B, even more preferably at least about 70-80%, 80-90%, 90-95% homologous to one of the sequences in Appendix B, and most preferably at least about 96%, 97%, 98%, or 99% homologous to one of the sequences in Appendix B.

To determine the percent homology of two amino acid sequences (*e.g.*, one of the sequences of Appendix B and a mutant form thereof) or of two nucleic acids, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in the sequence of one protein or nucleic acid for optimal alignment with the other protein or nucleic acid). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in one sequence (*e.g.*, one of the sequences of Appendix B) is occupied by the same amino acid residue or nucleotide as the corresponding position in the other sequence (*e.g.*, a mutant form of

09603208 062300

the sequence selected from Appendix B), then the molecules are homologous at that position (*i.e.*, as used herein amino acid or nucleic acid "homology" is equivalent to amino acid or nucleic acid "identity"). The percent homology between the two sequences is a function of the number of identical positions shared by the sequences (*i.e.*, % homology = # of identical positions/total # of positions x 100).

An isolated nucleic acid molecule encoding an SRT protein homologous to a protein sequence of Appendix B can be created by introducing one or more nucleotide substitutions, additions or deletions into a nucleotide sequence of Appendix A such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein. Mutations can be introduced into one of the sequences of Appendix A by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in an SRT protein is preferably replaced with another amino acid residue from the same side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of an SRT coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for an SRT activity described herein to identify mutants that retain SRT activity. Following mutagenesis of one of the sequences of Appendix A, the encoded protein can be expressed recombinantly and the activity of the protein can be determined using, for example, assays described herein (see Example 8 of the Exemplification).

In addition to the nucleic acid molecules encoding SRT proteins described above, another aspect of the invention pertains to isolated nucleic acid molecules which are antisense thereto. An "antisense" nucleic acid comprises a nucleotide sequence which is complementary to a "sense" nucleic acid encoding a protein, *e.g.*, complementary to the coding strand of a double-stranded DNA molecule or complementary to an mRNA sequence. Accordingly, an antisense nucleic acid can hydrogen bond to a sense nucleic acid. The antisense nucleic acid can be

complementary to an entire SRT coding strand, or to only a portion thereof. In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence encoding an SRT protein. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues (*e.g.*, the entire coding region of SEQ ID NO.: 120 (RXA00600) comprises nucleotides 1 to 1098). In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence encoding SRT. The term "noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (*i.e.*, also referred to as 5' and 3' untranslated regions).

Given the coding strand sequences encoding SRT disclosed herein (*e.g.*, the sequences set forth in Appendix A), antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of SRT mRNA, but more preferably is an oligonucleotide which is antisense to only a portion of the coding or noncoding region of SRT mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of SRT mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic acid of the invention can be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (*e.g.*, an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, *e.g.*, phosphorothioate derivatives and acridine substituted nucleotides can be used. Examples of modified nucleotides which can be used to generate the antisense nucleic acid include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-

amino-3-N-2-carboxypyril) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (*i.e.*, RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a cell or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding an SRT protein to thereby inhibit expression of the protein, *e.g.*, by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule which binds to DNA duplexes, through specific interactions in the major groove of the double helix. The antisense molecule can be modified such that it specifically binds to a receptor or an antigen expressed on a selected cell surface, *e.g.*, by linking the antisense nucleic acid molecule to a peptide or an antibody which binds to a cell surface receptor or antigen. The antisense nucleic acid molecule can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of the antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong prokaryotic, viral, or eukaryotic promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gaultier *et al.* (1987) *Nucleic Acids Res.* 15:6625-6641). The antisense nucleic acid molecule can also comprise a 2'-o-methylribonucleotide (Inoue *et al.* (1987) *Nucleic Acids Res.* 15:6131-6148) or a chimeric RNA-DNA analogue (Inoue *et al.* (1987) *FEBS Lett.* 215:327-330).

In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity which are capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they have a complementary region. Thus, ribozymes (*e.g.*, hammerhead ribozymes (described in Haselhoff and Gerlach (1988) *Nature* 334:585-591)) can be used to catalytically cleave SRT mRNA transcripts to thereby inhibit translation of SRT mRNA. A ribozyme having specificity for an SRT-encoding nucleic acid can be designed based upon the nucleotide sequence of an SRT cDNA disclosed herein (*i.e.*, SEQ ID NO:119 (RXA00600 in Appendix A)). For example, a derivative of a *Tetrahymena* L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in an SRT-encoding mRNA.

See, *e.g.*, Cech *et al.* U.S. Patent No. 4,987,071 and Cech *et al.* U.S. Patent No. 5,116,742. Alternatively, SRT mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, *e.g.*, Bartel, D. and Szostak, J.W. (1993) *Science* 261:1411-1418.

5 Alternatively, SRT gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of an SRT nucleotide sequence (*e.g.*, an SRT promoter and/or enhancers) to form triple helical structures that prevent transcription of an SRT gene in target cells. See generally, Helene, C. (1991) *Anticancer Drug Des.* 6(6):569-84; Helene, C. *et al.* (1992) *Ann. N.Y. Acad. Sci.* 660:27-10 36; and Maher, L.J. (1992) *Bioassays* 14(12):807-15.

B. Recombinant Expression Vectors and Host Cells

Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding an SRT protein (or a portion thereof). As
15 used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of
20 autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (*e.g.*, non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to
25 which they are operatively linked. Such vectors are referred to herein as "expression vectors". In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors,
30 such as viral vectors (*e.g.*, replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory
35 sequences, selected on the basis of the host cells to be used for expression, which is operatively linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, "operably linked" is intended to mean that the nucleotide sequence of

09603208 062300

interest is linked to the regulatory sequence(s) in a manner which allows for expression of the nucleotide sequence (*e.g.*, in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell). The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (*e.g.*, polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel; *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in many types of host cell and those which direct expression of the nucleotide sequence only in certain host cells.

Preferred regulatory sequences are, for example, promoters such as *cos*-, *tac*-, *trp*-, *tet*-, *trp-tet*-, *lpp*-, *lac*-, *lpp-lac*-, *lacI^q*-, *T7*-, *T5*-, *T3*-, *gal*-, *trc*-, *ara*-, *SP6*-, *arny*, *SPO2*, λ -*P_R*- or λ *P_L*, which are used preferably in bacteria. Additional regulatory sequences are, for example, promoters from yeasts and fungi, such as *ADC1*, *MF α* , *AC*, *P-60*, *CYC1*, *GAPDH*, *TEF*, *rp28*, *ADH*, promoters from plants such as *CaMV/35S*, *SSU*, *OCS*, *lib4*, *usp*, *STLS1*, *B33*, *nos* or *ubiquitin*- or *phaseolin*-promoters. It is also possible to use artificial promoters. It will be appreciated by one of ordinary skill in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. The expression vectors of the invention can be introduced into host cells to thereby produce proteins or peptides; including fusion proteins or peptides, encoded by nucleic acids as described herein (*e.g.*, *SRT* proteins, mutant forms of *SRT* proteins, fusion proteins, etc.).

The recombinant expression vectors of the invention can be designed for expression of *SRT* proteins in prokaryotic or eukaryotic cells. For example, *SRT* genes can be expressed in bacterial cells such as *C. glutamicum*, insect cells (using baculovirus expression vectors), yeast and other fungal cells (see Romanos, M.A. *et al.* (1992) "Foreign gene expression in yeast: a review", *Yeast* 8: 423-488; van den Hondel, C.A.M.J.J. *et al.* (1991) "Heterologous gene expression in filamentous fungi" in: *More Gene Manipulations in Fungi*, J.W. Bennet & L.L. Lasure, eds., p. 396-428: Academic Press: San Diego; and van den Hondel, C.A.M.J.J. & Punt, P.J. (1991) "Gene transfer systems and vector development for filamentous fungi, in: *Applied Molecular Genetics of Fungi*, Peberdy, J.F. *et al.*, eds., p. 1-28, Cambridge University Press: Cambridge), algae and multicellular plant cells (see Schmidt, R. and Willmitzer, L. (1988) High efficiency *Agrobacterium tumefaciens* -mediated transformation of *Arabidopsis thaliana* leaf and cotyledon explants" *Plant Cell Rep.*: 583-586), or mammalian cells. Suitable host cells are discussed further in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Alternatively, the

09603208 062300

recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

Expression of proteins in prokaryotes is most often carried out with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein. Such fusion vectors typically serve three purposes: 1) to increase expression of recombinant protein; 2) to increase the solubility of the recombinant protein; and 3) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase.

Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith, D.B. and Johnson, K.S. (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein. In one embodiment, the coding sequence of the SRT protein is cloned into a pGEX expression vector to create a vector encoding a fusion protein comprising, from the N-terminus to the C-terminus, GST-thrombin cleavage site-X protein. The fusion protein can be purified by affinity chromatography using glutathione-agarose resin. Recombinant SRT protein unfused to GST can be recovered by cleavage of the fusion protein with thrombin.

Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann *et al.*, (1988) *Gene* 69:301-315) pLG338, pACYC184, pBR322, pUC18, pUC19, pKC30, pRep4, pHS1, pHS2, pPLc236, pMBL24, pLG200, pUR290, pIN-III113-B1, λ gt11, pBdCl, and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 60-89; and Pouwels *et al.*, eds. (1985) *Cloning Vectors*. Elsevier: New York ISBN 0 444 904018). Target gene expression from the pTrc vector relies on host RNA polymerase transcription from a hybrid trp-lac fusion promoter. Target gene expression from the pET 11d vector relies on transcription from a T7 gn10-lac fusion promoter mediated by a coexpressed viral RNA polymerase (T7 gn1). This viral polymerase is supplied by host strains BL21(DE3) or HMS174(DE3) from a resident λ prophage harboring a T7 gn1 gene under the transcriptional control of the lacUV 5 promoter. For transformation of other varieties of bacteria, appropriate vectors may be selected. For example, the plasmids pIJ101, pIJ364, pIJ702 and pIJ361 are known to be useful in transforming

Streptomyces, while plasmids pUB110, pC194, or pBD214 are suited for transformation of Bacillus species. Several plasmids of use in the transfer of genetic information into Corynebacterium include pHM1519, pBL1, pSA77, or pAJ667 (Pouwels *et al.*, eds. (1985) Cloning Vectors. Elsevier: New York ISBN 0 444 904018)..

5 One strategy to maximize recombinant protein expression is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those
10 preferentially utilized in the bacterium chosen for expression, such as *C. glutamicum* (Wada *et al.* (1992) *Nucleic Acids Res.* 20:2111-2118). Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

In another embodiment, the SRT protein expression vector is a yeast expression vector. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari, *et al.*, (1987) *Embo J.* 6:229-234), 2 μ , pAG-1, Yep6, Yep13, pEMBLYe23, pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-943), pJRY88 (Schultz *et al.*, (1987) *Gene* 54:113-123), and pYES2 (Invitrogen Corporation, San Diego, CA). Vectors and methods for the construction of vectors appropriate for use in other fungi, such as the filamentous fungi, include those detailed in: van den Hondel, C.A.M.J.J. & Punt, P.J. (1991) "Gene transfer systems and vector development for filamentous fungi, in: Applied Molecular Genetics of Fungi, J.F. Peberdy, *et al.*, eds., p. 1-28, Cambridge University Press: Cambridge, and Pouwels *et al.*, eds. (1985) Cloning Vectors. Elsevier: New York (IBSN 0 444 904018).

25 Alternatively, the SRT proteins of the invention can be expressed in insect cells using baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (*e.g.*, Sf 9 cells) include the pAc series (Smith *et al.* (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow and Summers (1989) *Virology* 170:31-39).

30 In another embodiment, the SRT proteins of the invention may be expressed in unicellular plant cells (such as algae) or in plant cells from higher plants (*e.g.*, the spermatophytes, such as crop plants). Examples of plant expression vectors include those detailed in: Becker, D., Kemper, E., Schell, J. and Masterson, R. (1992) "New plant binary vectors with selectable markers located proximal to the left border", *Plant*
35 *Mol. Biol.* 20: 1195-1197; and Bevan, M.W. (1984) "Binary *Agrobacterium* vectors for plant transformation", *Nucl. Acid. Res.* 12: 8711-8721, and include pLGV23, pGHIac+,

In yet another embodiment, a nucleic acid of the invention is expressed in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, B. (1987) *Nature* 329:840) and pMT2PC (Kaufman *et al.* (1987) *EMBO J.* 6:187-195). When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40. For other suitable expression systems for both prokaryotic and eukaryotic cells see chapters 16 and 17 of Sambrook, J., Fritsch, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (e.g., tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert *et al.* (1987) *Genes Dev.* 1:268-277), lymphoid-specific promoters (Calame and Eaton (1988) *Adv. Immunol.* 43:235-275), in particular promoters of T cell receptors (Winoto and Baltimore (1989) *EMBO J.* 8:729-733) and immunoglobulins (Banerji *et al.* (1983) *Cell* 33:729-740; Queen and Baltimore (1983) *Cell* 33:741-748), neuron-specific promoters (e.g., the neurofilament promoter; Byrne and Ruddle (1989) *PNAS* 86:5473-5477), pancreas-specific promoters (Edlund *et al.* (1985) *Science* 230:912-916), and mammary gland-specific promoters (e.g., milk whey promoter; U.S. Patent No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, for example the murine hox promoters (Kessel and Gruss (1990) *Science* 249:374-379) and the α -fetoprotein promoter (Campes and Tilghman (1989) *Genes Dev.* 3:537-546).

30 The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an antisense orientation. That is, the DNA molecule is operatively linked to a regulatory sequence in a manner which allows for expression (by transcription of the DNA molecule) of an RNA molecule which is antisense to SRT mRNA. Regulatory sequences operatively
35 linked to a nucleic acid cloned in the antisense orientation can be chosen which direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be chosen which

direct constitutive, tissue specific or cell type specific expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense nucleic acids are produced under the control of a high efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced. For a discussion of the regulation of gene expression using antisense genes see Weintraub, H. *et al.*, Antisense RNA as a molecular tool for genetic analysis, *Reviews - Trends in Genetics*, Vol. 1(1) 1986.

Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

A host cell can be any prokaryotic or eukaryotic cell. For example, an SRT protein can be expressed in bacterial cells such as *C. glutamicum*, insect cells, yeast or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those of ordinary skill in the art. Microorganisms related to *Corynebacterium glutamicum* which may be conveniently used as host cells for the nucleic acid and protein molecules of the invention are set forth in Table 3.

Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (*e.g.*, linear DNA or RNA (*e.g.*, a linearized vector or a gene construct alone without a vector) or nucleic acid in the form of a vector (*e.g.*, a plasmid, phage, phasmid, phagemid, transposon or other DNA)) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, *et al.* (*Molecular Cloning: A Laboratory Manual*, 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989), and other laboratory manuals.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (*e.g.*, resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred

00603208-062300

5

10

15

30

35

deletions, or inversions, but still encodes a functional SRT protein. In still another embodiment, one or more of the regulatory regions (*e.g.*, a promoter, repressor, or inducer) of an SRT gene in a microorganism has been altered (*e.g.*, by deletion, truncation, inversion, or point mutation) such that the expression of the SRT gene is modulated. One of ordinary skill in the art will appreciate that host cells containing more than one of the described SRT gene and protein modifications may be readily produced using the methods of the invention, and are meant to be included in the present invention.

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce (*i.e.*, express) an SRT protein. Accordingly, the invention further provides methods for producing SRT proteins using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention (into which a recombinant expression vector encoding an SRT protein has been introduced, or into which genome has been introduced a gene encoding a wild-type or altered SRT protein) in a suitable medium until SRT protein is produced. In another embodiment, the method further comprises isolating SRT proteins from the medium or the host cell.

C. Isolated SRT Proteins

Another aspect of the invention pertains to isolated SRT proteins, and biologically active portions thereof. An "isolated" or "purified" protein or biologically active portion thereof is substantially free of cellular material when produced by recombinant DNA techniques, or chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of SRT protein in which the protein is separated from cellular components of the cells in which it is naturally or recombinantly produced. In one embodiment, the language "substantially free of cellular material" includes preparations of SRT protein having less than about 30% (by dry weight) of non-SRT protein (also referred to herein as a "contaminating protein"), more preferably less than about 20% of non-SRT protein, still more preferably less than about 10% of non-SRT protein, and most preferably less than about 5% non-SRT protein. When the SRT protein or biologically active portion thereof is recombinantly produced, it is also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation. The language "substantially free of chemical precursors or other chemicals" includes preparations of SRT protein in which the protein is separated from chemical precursors or other chemicals which are involved in the synthesis of the

protein. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of SRT protein having less than about 30% (by dry weight) of chemical precursors or non-SRT chemicals, more preferably less than about 20% chemical precursors or non-SRT chemicals, still more preferably less than about 10% chemical precursors or non-SRT chemicals, and most preferably less than about 5% chemical precursors or non-SRT chemicals. In preferred embodiments, isolated proteins or biologically active portions thereof lack contaminating proteins from the same organism from which the SRT protein is derived. Typically, such proteins are produced by recombinant expression of, for example, a *C. glutamicum* SRT protein in a microorganism such as *C. glutamicum*.

An isolated SRT protein or a portion thereof of the invention can contribute to the resistance or tolerance of *C. glutamicum* to one or more chemical or environmental stresses or hazards, or has one or more of the activities set forth in Table 1. In preferred embodiments, the protein or portion thereof comprises an amino acid sequence which is sufficiently homologous to an amino acid sequence of Appendix B such that the protein or portion thereof maintains the ability to mediate the resistance or tolerance of *C. glutamicum* to one or more chemical or environmental stresses or hazards. The portion of the protein is preferably a biologically active portion as described herein. In another preferred embodiment, an SRT protein of the invention has an amino acid sequence shown in Appendix B. In yet another preferred embodiment, the SRT protein has an amino acid sequence which is encoded by a nucleotide sequence which hybridizes, *e.g.*, hybridizes under stringent conditions, to a nucleotide sequence of Appendix A. In still another preferred embodiment, the SRT protein has an amino acid sequence which is encoded by a nucleotide sequence that is at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, or 60%, preferably at least about 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, or 70%, more preferably at least about 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, or 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, or 90%, or 91%, 92%, 93%, 94%, and even more preferably at least about 95%, 96%, 97%, 98%, 99% or more homologous to one of the nucleic acid sequences of Appendix A, or a portion thereof. Ranges and identity values intermediate to the above-recited values, (*e.g.*, 70-90% identical or 80-95% identical) are also intended to be encompassed by the present invention. For example, ranges of identity values using a combination of any of the above values recited as upper and/or lower limits are intended to be included. The preferred SRT proteins of the present invention also preferably possess at least one of the SRT activities described herein. For example, a preferred SRT protein of the present invention includes an amino acid sequence encoded by a nucleotide sequence which hybridizes, *e.g.*, hybridizes under stringent conditions, to a

00603208.062300

nucleotide sequence of Appendix A, and which can increase the resistance or tolerance of *C. glutamicum* to one or more environmental or chemical stresses, or which has one or more of the activities set forth in Table 1.

In other embodiments, the SRT protein is substantially homologous to an amino acid sequence of Appendix B and retains the functional activity of the protein of one of the sequences of Appendix B yet differs in amino acid sequence due to natural variation or mutagenesis, as described in detail in subsection I above. Accordingly, in another embodiment, the SRT protein is a protein which comprises an amino acid sequence which is at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, or 60%, preferably at least about 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, or 70%, more preferably at least about 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, or 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, or 90%, or 91%, 92%, 93%, 94%, and even more preferably at least about 95%, 96%, 97%, 98%, 99% or more homologous to an entire amino acid sequence of Appendix B and which has at least one of the SRT activities described herein. Ranges and identity values intermediate to the above-recited values, (e.g., 70-90% identical or 80-95% identical) are also intended to be encompassed by the present invention. For example, ranges of identity values using a combination of any of the above values recited as upper and/or lower limits are intended to be included. In another embodiment, the invention pertains to a full length *C. glutamicum* protein which is substantially homologous to an entire amino acid sequence of Appendix B.

Biologically active portions of an SRT protein include peptides comprising amino acid sequences derived from the amino acid sequence of an SRT protein, e.g., an amino acid sequence shown in Appendix B or the amino acid sequence of a protein homologous to an SRT protein, which include fewer amino acids than a full length SRT protein or the full length protein which is homologous to an SRT protein, and exhibit at least one activity of an SRT protein. Typically, biologically active portions (peptides, e.g., peptides which are, for example, 5, 10, 15, 20, 30, 35, 36, 37, 38, 39, 40, 50, 100 or more amino acids in length) comprise a domain or motif with at least one activity of an SRT protein. Moreover, other biologically active portions, in which other regions of the protein are deleted, can be prepared by recombinant techniques and evaluated for one or more of the activities described herein. Preferably, the biologically active portions of an SRT protein include one or more selected domains/motifs or portions thereof having biological activity.

SRT proteins are preferably produced by recombinant DNA techniques. For example, a nucleic acid molecule encoding the protein is cloned into an expression vector (as described above), the expression vector is introduced into a host cell (as

09603208-062300

described above) and the SRT protein is expressed in the host cell. The SRT protein can then be isolated from the cells by an appropriate purification scheme using standard protein purification techniques. Alternative to recombinant expression, an SRT protein, polypeptide, or peptide can be synthesized chemically using standard peptide synthesis techniques. Moreover, native SRT protein can be isolated from cells (e.g., endothelial cells), for example using an anti-SRT antibody, which can be produced by standard techniques utilizing an SRT protein or fragment thereof of this invention.

The invention also provides SRT chimeric or fusion proteins. As used herein, an SRT "chimeric protein" or "fusion protein" comprises an SRT polypeptide operatively linked to a non-SRT polypeptide. An "SRT polypeptide" refers to a polypeptide having an amino acid sequence corresponding to SRT, whereas a "non-SRT polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein which is not substantially homologous to the SRT protein, e.g., a protein which is different from the SRT protein and which is derived from the same or a different organism. Within the fusion protein, the term "operatively linked" is intended to indicate that the SRT polypeptide and the non-SRT polypeptide are fused in-frame to each other. The non-SRT polypeptide can be fused to the N-terminus or C-terminus of the SRT polypeptide. For example, in one embodiment the fusion protein is a GST-SRT fusion protein in which the SRT sequences are fused to the C-terminus of the GST sequences. Such fusion proteins can facilitate the purification of recombinant SRT proteins. In another embodiment, the fusion protein is an SRT protein containing a heterologous signal sequence at its N-terminus. In certain host cells (e.g., mammalian host cells), expression and/or secretion of an SRT protein can be increased through use of a heterologous signal sequence.

Preferably, an SRT chimeric or fusion protein of the invention is produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, for example by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel et al. John Wiley & Sons: 1992). Moreover, many expression vectors are commercially

available that already encode a fusion moiety (e.g., a GST polypeptide). An SRT-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the SRT protein.

Homologues of the SRT protein can be generated by mutagenesis, e.g., discrete point mutation or truncation of the SRT protein. As used herein, the term "homologue" refers to a variant form of the SRT protein which acts as an agonist or antagonist of the activity of the SRT protein. An agonist of the SRT protein can retain substantially the same, or a subset, of the biological activities of the SRT protein. An antagonist of the SRT protein can inhibit one or more of the activities of the naturally occurring form of the SRT protein, by, for example, competitively binding to a downstream or upstream member of the SRT system which includes the SRT protein. Thus, the *C. glutamicum* SRT protein and homologues thereof of the present invention may increase the tolerance or resistance of *C. glutamicum* to one or more chemical or environmental stresses.

In an alternative embodiment, homologues of the SRT protein can be identified by screening combinatorial libraries of mutants, e.g., truncation mutants, of the SRT protein for SRT protein agonist or antagonist activity. In one embodiment, a variegated library of SRT variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library of SRT variants can be produced by, for example, enzymatically ligating a mixture of synthetic oligonucleotides into gene sequences such that a degenerate set of potential SRT sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (e.g., for phage display) containing the set of SRT sequences therein. There are a variety of methods which can be used to produce libraries of potential SRT homologues from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can be performed in an automatic DNA synthesizer, and the synthetic gene then ligated into an appropriate expression vector. Use of a degenerate set of genes allows for the provision, in one mixture, of all of the sequences encoding the desired set of potential SRT sequences. Methods for synthesizing degenerate oligonucleotides are known in the art (see, e.g., Narang, S.A. (1983) *Tetrahedron* 39:3; Itakura *et al.* (1984) *Annu. Rev. Biochem.* 53:323; Itakura *et al.* (1984) *Science* 198:1056; Ike *et al.* (1983) *Nucleic Acid Res.* 11:477).

In addition, libraries of fragments of the SRT protein coding can be used to generate a variegated population of SRT fragments for screening and subsequent selection of homologues of an SRT protein. In one embodiment, a library of coding sequence fragments can be generated by treating a double stranded PCR fragment of an SRT coding sequence with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double stranded DNA, renaturing the DNA to

09603208-062300

form double stranded DNA which can include sense/antisense pairs from different nicked products, removing single stranded portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, an expression library can be derived which encodes N-terminal, C-terminal and internal fragments of various sizes of the SRT protein.

Several techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property. Such techniques are adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of SRT homologues. The most widely used techniques, which are amenable to high through-put analysis, for screening large gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a new technique which enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify SRT homologues (Arkin and Yourvan (1992) *PNAS* 89:7811-7815; Delgrave *et al.* (1993) *Protein Engineering* 6(3):327-331).

In another embodiment, cell based assays can be exploited to analyze a variegated SRT library, using methods well known in the art.

D. Uses and Methods of the Invention

The nucleic acid molecules, proteins, protein homologues, fusion proteins, primers, vectors, and host cells described herein can be used in one or more of the following methods: identification of *C. glutamicum* and related organisms; mapping of genomes of organisms related to *C. glutamicum*; identification and localization of *C. glutamicum* sequences of interest; evolutionary studies; determination of SRT protein regions required for function; modulation of an SRT protein activity; modulation of the activity of an SRT pathway; and modulation of cellular production of a desired compound, such as a fine chemical.

The SRT nucleic acid molecules of the invention have a variety of uses. First, they may be used to identify an organism as being *Corynebacterium glutamicum* or a close relative thereof. Also, they may be used to identify the presence of *C. glutamicum* or a relative thereof in a mixed population of microorganisms. The invention provides the nucleic acid sequences of a number of *C. glutamicum* genes; by probing the extracted genomic DNA of a culture of a unique or mixed population of microorganisms

under stringent conditions with a probe spanning a region of a *C. glutamicum* gene which is unique to this organism, one can ascertain whether this organism is present.

Although *Corynebacterium glutamicum* itself is nonpathogenic, it is related to pathogenic species, such as *Corynebacterium diphtheriae*. *Corynebacterium diphtheriae* is the causative agent of diphtheria, a rapidly developing, acute, febrile infection which involves both local and systemic pathology. In this disease, a local lesion develops in the upper respiratory tract and involves necrotic injury to epithelial cells; the bacilli secrete toxin which is disseminated through this lesion to distal susceptible tissues of the body. Degenerative changes brought about by the inhibition of protein synthesis in these tissues, which include heart, muscle, peripheral nerves, adrenals, kidneys, liver and spleen, result in the systemic pathology of the disease. Diphtheria continues to have high incidence in many parts of the world, including Africa, Asia, Eastern Europe and the independent states of the former Soviet Union. An ongoing epidemic of diphtheria in the latter two regions has resulted in at least 5,000 deaths since 1990.

In one embodiment, the invention provides a method of identifying the presence or activity of *Corynebacterium diphtheriae* in a subject. This method includes detection of one or more of the nucleic acid or amino acid sequences of the invention (*e.g.*, the sequences set forth in Appendix A or Appendix B) in a subject, thereby detecting the presence or activity of *Corynebacterium diphtheriae* in the subject. *C. glutamicum* and *C. diphtheriae* are related bacteria, and many of the nucleic acid and protein molecules in *C. glutamicum* are homologous to *C. diphtheriae* nucleic acid and protein molecules, and can therefore be used to detect *C. diphtheriae* in a subject.

The nucleic acid and protein molecules of the invention may also serve as markers for specific regions of the genome. This has utility not only in the mapping of the genome, but also for functional studies of *C. glutamicum* proteins. For example, to identify the region of the genome to which a particular *C. glutamicum* DNA-binding protein binds, the *C. glutamicum* genome could be digested, and the fragments incubated with the DNA-binding protein. Those which bind the protein may be additionally probed with the nucleic acid molecules of the invention, preferably with readily detectable labels; binding of such a nucleic acid molecule to the genome fragment enables the localization of the fragment to the genome map of *C. glutamicum*, and, when performed multiple times with different enzymes, facilitates a rapid determination of the nucleic acid sequence to which the protein binds. Further, the nucleic acid molecules of the invention may be sufficiently homologous to the sequences of related species such that these nucleic acid molecules may serve as markers for the construction of a genomic map in related bacteria, such as *Brevibacterium lactofermentum*.

09603208 "062300

09603208-062300

The SRT nucleic acid molecules of the invention are also useful for evolutionary and protein structural studies. The resistance processes in which the molecules of the invention participate are utilized by a wide variety of cells; by comparing the sequences of the nucleic acid molecules of the present invention to those encoding similar enzymes from other organisms, the evolutionary relatedness of the organisms can be assessed. Similarly, such a comparison permits an assessment of which regions of the sequence are conserved and which are not, which may aid in determining those regions of the protein which are essential for the functioning of the enzyme. This type of determination is of value for protein engineering studies and may give an indication of what the protein can tolerate in terms of mutagenesis without losing function.

The genes of the invention, *e.g.*, the gene encoding LMRB (SEQ ID NO:1) or other gene of the invention encoding a chemical or environmental resistance or tolerance protein (*e.g.*, resistance against one or more antibiotics), may be used as genetic markers for the genetic transformation of (*e.g.*, the transfer of additional genes into or disruption of preexisting genes of) organisms such as *C. glutamicum* or other bacterial species. Use of these nucleic acid molecules permits efficient selection of organisms which have incorporated a given transgene cassette (*e.g.*, a plasmid, phage, phasmid, phagemid, transposon, or other nucleic acid element), based on a trait which permits the survival of the organism in an otherwise hostile or toxic environment (*e.g.*, in the presence of an antimicrobial compound). By employing one or more of the genes of the invention as genetic markers, the speed and ease with which organisms having desirable transformed traits (*e.g.*, modulated fine chemical production) are engineered and isolated are improved. While it is advantageous to use the genes of the invention for selection of transformed *C. glutamicum* and related bacteria, it is possible, as described herein, to use homologs (*e.g.*, homologs from other organisms), allelic variants or fragments of the gene retaining desired activity. Furthermore, 5' and 3' regulatory elements of the genes of the invention may be modified as described herein (*e.g.*, by nucleotide substitution, insertion, deletion, or replacement with a more desirable genetic element) to modulate the transcription of the gene. For example, an LMRB variant in which the nucleotide sequence in the region from -1 to -200 5' to the start codon has been altered to modulate (preferably increase) the transcription and/or translation of LMRB may be employed, as can constructs in which a gene of the invention (*e.g.*, the LMRB gene (SEQ ID NO:1)) is functionally coupled to one or more regulatory signals (*e.g.*, inducer or repressor binding sequences) which can be used for modulating gene expression. Similarly, more than one copy of a gene (functional or inactivated) of the invention may be employed.

An additional application of the genes of the invention (*e.g.*, the gene encoding LMRB (SEQ ID NO:1) or other drug- or antibiotic-resistance gene) is in the discovery of new antibiotics which are active against *Corynebacteria* and/or other bacteria. For example, a gene of the invention may be expressed (or overexpressed) in a suitable host
5 to generate an organism with increased resistance to one or more drugs or antibiotics (in the case of LMRB, lincosamides in particular, especially lincomycin). This type of resistant host can subsequently be used to screen for chemicals with bacteriostatic and/or bacteriocidal activity, such as novel antibiotic compounds. It is possible, in particular, to use the genes of the invention (*e.g.*, the LMRB gene) to identify new antibiotics
10 which are active against those microorganisms which are already resistant to standard antibiotic compounds.

The invention provides methods for screening molecules which modulate the activity of an SRT protein, either by interacting with the protein itself or a substrate or binding partner of the SRT protein, or by modulating the transcription or translation of a
15 SRT nucleic acid molecule of the invention. In such methods, a microorganism expressing one or more SRT proteins of the invention is contacted with one or more test compounds, and the effect of each test compound on the activity or level of expression of the SRT protein is assessed.

Manipulation of the SRT nucleic acid molecules of the invention may result in
20 the production of SRT proteins having functional differences from the wild-type SRT proteins. These proteins may be improved in efficiency or activity, may be present in greater numbers in the cell than is usual, or may be decreased in efficiency or activity. The goal of such manipulations is to increase the viability and activity of the cell when the cell is exposed to the environmental and chemical stresses and hazards which
25 frequently accompany large-scale fermentative culture. Thus, by increasing the activity or copy number of a heat-shock-regulated protease, one may increase the ability of the cell to destroy incorrectly folded proteins, which may otherwise interfere with normal cellular functioning (for example, by continuing to bind substrates or cofactors although the protein lacks the activity to act on these molecules appropriately). The same is true
30 for the overexpression or optimization of activity of one or more chaperone molecules induced by heat or cold shock. These proteins aid in the correct folding of nascent polypeptide chains, and thus their increased activity or presence should increase the percentage of correctly folded proteins in the cell, which in turn should increase the overall metabolic efficiency and viability of the cells in culture. The overexpression or
35 optimization of the transporter molecules activated by osmotic shock should result in an increased ability on the part of the cell to maintain intracellular homeostasis, thereby increasing the viability of these cells in culture. Similarly, the overproduction or

09603208-062300

increase in activity by mutagenesis of proteins involved in the development of cellular resistance to chemical stresses of various kinds (either by transport of the offending chemical out of the cell or by modification of the chemical to a less hazardous substance) should increase the fitness of the organism in the environment containing the hazardous substance (*i.e.*, large-scale fermentative culture), and thereby may permit relatively larger numbers of cells to survive in such a culture. The net effect of all of these mutagenesis strategies is to increase the quantity of fine-chemical-producing compounds in the culture, thereby increasing the yield, production, and/or efficiency of production of one or more desired fine chemicals from the culture.

10 This aforementioned list of mutagenesis strategies for SRT proteins to result in increased yields of a desired compound is not meant to be limiting; variations on these mutagenesis strategies will be readily apparent to one of ordinary skill in the art. By these mechanisms, the nucleic acid and protein molecules of the invention may be utilized to generate *C. glutamicum* or related strains of bacteria expressing mutated SRT
15 nucleic acid and protein molecules such that the yield, production, and/or efficiency of production of a desired compound is improved. This desired compound may be any natural product of *C. glutamicum*, which includes the final products of biosynthesis pathways and intermediates of naturally-occurring metabolic pathways, as well as molecules which do not naturally occur in the metabolism of *C. glutamicum*, but which
20 are produced by a *C. glutamicum* strain of the invention.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patent applications, patents, published patent applications, Tables, Appendices, and the sequence listing cited throughout this application are hereby incorporated by reference.

25

Exemplification

Example 1: Preparation of total genomic DNA of *Corynebacterium glutamicum* ATCC 13032

30 A culture of *Corynebacterium glutamicum* (ATCC 13032) was grown overnight at 30°C with vigorous shaking in BHI medium (Difco). The cells were harvested by centrifugation, the supernatant was discarded and the cells were resuspended in 5 ml buffer-I (5% of the original volume of the culture — all indicated volumes have been calculated for 100 ml of culture volume). Composition of buffer-I: 140.34 g/l sucrose,
35 2.46 g/l MgSO₄ x 7H₂O, 10 ml/l KH₂PO₄ solution (100 g/l, adjusted to pH 6.7 with KOH), 50 ml/l M12 concentrate (10 g/l (NH₄)₂SO₄, 1 g/l NaCl, 2 g/l MgSO₄ x 7H₂O, 0.2 g/l CaCl₂, 0.5 g/l yeast extract (Difco), 10 ml/l trace-elements-mix (200 mg/l FeSO₄

09603208-062300

09603208-062300
x H₂O, 10 mg/l ZnSO₄ x 7 H₂O, 3 mg/l MnCl₂ x 4 H₂O, 30 mg/l H₃BO₃, 20 mg/l CoCl₂ x 6 H₂O, 1 mg/l NiCl₂ x 6 H₂O, 3 mg/l Na₂MoO₄ x 2 H₂O, 500 mg/l complexing agent (EDTA or citric acid), 100 ml/l vitamins-mix (0.2 mg/l biotin, 0.2 mg/l folic acid, 20 mg/l p-amino benzoic acid, 20 mg/l riboflavin, 40 mg/l ca-panthothenate, 140 mg/l
5 nicotinic acid, 40 mg/l pyridoxole hydrochloride, 200 mg/l myo-inositol). Lysozyme was added to the suspension to a final concentration of 2.5 mg/ml. After an approximately 4 h incubation at 37°C, the cell wall was degraded and the resulting protoplasts are harvested by centrifugation. The pellet was washed once with 5 ml buffer-I and once with 5 ml TE-buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8). The
10 pellet was resuspended in 4 ml TE-buffer and 0.5 ml SDS solution (10%) and 0.5 ml NaCl solution (5 M) are added. After adding of proteinase K to a final concentration of 200 µg/ml, the suspension is incubated for ca.18 h at 37°C. The DNA was purified by extraction with phenol, phenol-chloroform-isoamylalcohol and chloroform-isoamylalcohol using standard procedures. Then, the DNA was precipitated by adding
15 1/50 volume of 3 M sodium acetate and 2 volumes of ethanol, followed by a 30 min incubation at -20°C and a 30 min centrifugation at 12,000 rpm in a high speed centrifuge using a SS34 rotor (Sorvall). The DNA was dissolved in 1 ml TE-buffer containing 20 µg/ml RNaseA and dialysed at 4°C against 1000 ml TE-buffer for at least 3 hours. During this time, the buffer was exchanged 3 times. To aliquots of 0.4 ml of the
20 dialysed DNA solution, 0.4 ml of 2 M LiCl and 0.8 ml of ethanol are added. After a 30 min incubation at -20°C, the DNA was collected by centrifugation (13,000 rpm, Biofuge Fresco, Heraeus, Hanau, Germany). The DNA pellet was dissolved in TE-buffer. DNA prepared by this procedure could be used for all purposes, including southern blotting or construction of genomic libraries.

25

Example 2: Construction of genomic libraries in *Escherichia coli* of *Corynebacterium glutamicum* ATCC13032.

Using DNA prepared as described in Example 1, cosmid and plasmid libraries were constructed according to known and well established methods (*see e.g.*, Sambrook, J. *et al.*
30 (1989) "Molecular Cloning : A Laboratory Manual", Cold Spring Harbor Laboratory Press, or Ausubel, F.M. *et al.* (1994) "Current Protocols in Molecular Biology", John Wiley & Sons.)

Any plasmid or cosmid could be used. Of particular use were the plasmids pBR322 (Sutcliffe, J.G. (1979) *Proc. Natl. Acad. Sci. USA*, 75:3737-3741); pACYC177 (Change &
35 Cohen (1978) *J. Bacteriol* 134:1141-1156), plasmids of the pBS series (pBSSK+, pBSSK- and others; Stratagene, LaJolla, USA), or cosmids as SuperCos1 (Stratagene, LaJolla, USA) or Lorist6 (Gibson, T.J., Rosenthal A. and Waterson, R.H. (1987) *Gene* 53:283-286. Gene libraries

specifically for use in *C. glutamicum* may be constructed using plasmid pSL109 (Lee, H.-S. and A. J. Sinskey (1994) *J. Microbiol. Biotechnol.* 4: 256-263).

Example 3: DNA Sequencing and Computational Functional Analysis

5 Genomic libraries as described in Example 2 were used for DNA sequencing according to standard methods, in particular by the chain termination method using ABI377 sequencing machines (see *e.g.*, Fleischman, R.D. *et al.* (1995) "Whole-genome Random Sequencing and Assembly of Haemophilus Influenzae Rd., *Science*, 269:496-512). Sequencing primers with the following nucleotide sequences were used: 5'-
10 GGAAACAGTATGACCATG-3' or 5'-GTAAAACGACGGCCAGT-3'.

Example 4: *In vivo* Mutagenesis

15 *In vivo* mutagenesis of *Corynebacterium glutamicum* can be performed by passage of plasmid (or other vector) DNA through *E. coli* or other microorganisms (*e.g.* *Bacillus* spp. or yeasts such as *Saccharomyces cerevisiae*) which are impaired in their capabilities to maintain the integrity of their genetic information. Typical mutator strains have mutations in the genes for the DNA repair system (*e.g.*, mutHLS, mutD, mutT, etc.; for reference, see Rupp, W.D. (1996) DNA repair mechanisms, in: *Escherichia coli* and *Salmonella*, p. 2277-2294, ASM: Washington.) Such strains are well known to those of ordinary skill in the art. The use of such
20 strains is illustrated, for example, in Greener, A. and Callahan, M. (1994) *Strategies* 7: 32-34.

Example 5: DNA Transfer Between *Escherichia coli* and *Corynebacterium glutamicum*

25 Several *Corynebacterium* and *Brevibacterium* species contain endogenous plasmids (as *e.g.*, pHM1519 or pBL1) which replicate autonomously (for review see, *e.g.*, Martin, J.F. *et al.* (1987) *Biotechnology*, 5:137-146). Shuttle vectors for *Escherichia coli* and *Corynebacterium glutamicum* can be readily constructed by using standard vectors for *E. coli* (Sambrook, J. *et al.* (1989), "Molecular Cloning: A Laboratory Manual", Cold Spring Harbor Laboratory Press or Ausubel, F.M. *et al.* (1994) "Current Protocols in
30 Molecular Biology", John Wiley & Sons) to which a origin or replication for and a suitable marker from *Corynebacterium glutamicum* is added. Such origins of replication are preferably taken from endogenous plasmids isolated from *Corynebacterium* and *Brevibacterium* species. Of particular use as transformation markers for these species are genes for kanamycin resistance (such as those derived from the Tn5 or Tn903
35 transposons) or chloramphenicol (Winnacker, E.L. (1987) "From Genes to Clones — Introduction to Gene Technology, VCH, Weinheim). There are numerous examples in the literature of the construction of a wide variety of shuttle vectors which replicate in both *E.*

coli and *C. glutamicum*, and which can be used for several purposes, including gene over-expression (for reference, see *e.g.*, Yoshihama, M. *et al.* (1985) *J. Bacteriol.* 162:591-597, Martin J.F. *et al.* (1987) *Biotechnology*, 5:137-146 and Eikmanns, B.J. *et al.* (1991) *Gene*, 102:93-98).

5 Using standard methods, it is possible to clone a gene of interest into one of the shuttle vectors described above and to introduce such a hybrid vector into strains of *Corynebacterium glutamicum*. Transformation of *C. glutamicum* can be achieved by protoplast transformation (Kastsumata, R. *et al.* (1984) *J. Bacteriol.* 159:306-311), electroporation (Liebl, E. *et al.* (1989) *FEMS Microbiol. Letters*, 53:399-303) and in cases
10 where special vectors are used, also by conjugation (as described *e.g.* in Schäfer, A *et al.* (1990) *J. Bacteriol.* 172:1663-1666). It is also possible to transfer the shuttle vectors for *C. glutamicum* to *E. coli* by preparing plasmid DNA from *C. glutamicum* (using standard methods well-known in the art) and transforming it into *E. coli*. This transformation step can be performed using standard methods, but it is advantageous to use an Mcr-deficient
15 *E. coli* strain, such as NM522 (Gough & Murray (1983) *J. Mol. Biol.* 166:1-19).

 Genes may be overexpressed in *C. glutamicum* strains using plasmids which comprise pCG1 (U.S. Patent No. 4,617,267) or fragments thereof, and optionally the gene for kanamycin resistance from TN903 (Grindley, N.D. and Joyce, C.M. (1980) *Proc. Natl. Acad. Sci. USA* 77(12): 7176-7180). In addition, genes may be
20 overexpressed in *C. glutamicum* strains using plasmid pSL109 (Lee, H.-S. and A. J. Sinskey (1994) *J. Microbiol. Biotechnol.* 4: 256-263).

 Aside from the use of replicative plasmids, gene overexpression can also be achieved by integration into the genome. Genomic integration in *C. glutamicum* or other *Corynebacterium* or *Brevibacterium* species may be accomplished by well-known
25 methods, such as homologous recombination with genomic region(s), restriction endonuclease mediated integration (REMI) (see, *e.g.*, DE Patent 19823834), or through the use of transposons. It is also possible to modulate the activity of a gene of interest by modifying the regulatory regions (*e.g.*, a promoter, a repressor, and/or an enhancer) by sequence modification, insertion, or deletion using site-directed methods (such as
30 homologous recombination) or methods based on random events (such as transposon mutagenesis or REMI). Nucleic acid sequences which function as transcriptional terminators may also be inserted 3' to the coding region of one or more genes of the invention; such terminators are well-known in the art and are described, for example, in Winnacker, E.L. (1987) *From Genes to Clones – Introduction to Gene Technology*. VCH:
35 Weinheim.

09603208-062300

Example 6: Assessment of the Expression of the Mutant Protein

Observations of the activity of a mutated protein in a transformed host cell rely on the fact that the mutant protein is expressed in a similar fashion and in a similar quantity to that of the wild-type protein. A useful method to ascertain the level of transcription of the mutant gene (an indicator of the amount of mRNA available for translation to the gene product) is to perform a Northern blot (for reference see, for example, Ausubel *et al.* (1988) *Current Protocols in Molecular Biology*, Wiley: New York), in which a primer designed to bind to the gene of interest is labeled with a detectable tag (usually radioactive or chemiluminescent), such that when the total RNA of a culture of the organism is extracted, run on gel, transferred to a stable matrix and incubated with this probe, the binding and quantity of binding of the probe indicates the presence and also the quantity of mRNA for this gene. This information is evidence of the degree of transcription of the mutant gene. Total cellular RNA can be prepared from *Corynebacterium glutamicum* by several methods, all well-known in the art, such as that described in Bormann, E.R. *et al.* (1992) *Mol. Microbiol.* 6: 317-326.

To assess the presence or relative quantity of protein translated from this mRNA, standard techniques, such as a Western blot, may be employed (see, for example, Ausubel *et al.* (1988) *Current Protocols in Molecular Biology*, Wiley: New York). In this process, total cellular proteins are extracted, separated by gel electrophoresis, transferred to a matrix such as nitrocellulose, and incubated with a probe, such as an antibody, which specifically binds to the desired protein. This probe is generally tagged with a chemiluminescent or colorimetric label which may be readily detected. The presence and quantity of label observed indicates the presence and quantity of the desired mutant protein present in the cell.

Example 7: Growth of Genetically Modified *Corynebacterium glutamicum* — Media and Culture Conditions

Genetically modified *Corynebacteria* are cultured in synthetic or natural growth media. A number of different growth media for *Corynebacteria* are both well-known and readily available (Lieb *et al.* (1989) *Appl. Microbiol. Biotechnol.*, 32:205-210; von der Osten *et al.* (1998) *Biotechnology Letters*, 11:11-16; Patent DE 4,120,867; Liebl (1992) "The Genus *Corynebacterium*, in: *The Prokaryotes*, Volume II, Balows, A. *et al.*, eds. Springer-Verlag). These media consist of one or more carbon sources, nitrogen sources, inorganic salts, vitamins and trace elements. Preferred carbon sources are sugars, such as mono-, di-, or polysaccharides. For example, glucose, fructose, mannose, galactose, ribose, sorbose, ribulose, lactose, maltose, sucrose, raffinose, starch or cellulose serve as very good carbon sources. It is also possible to supply sugar to the media via complex

00603208-06300
compounds such as molasses or other by-products from sugar refinement. It can also be advantageous to supply mixtures of different carbon sources. Other possible carbon sources are alcohols and organic acids, such as methanol, ethanol, acetic acid or lactic acid. Nitrogen sources are usually organic or inorganic nitrogen compounds, or materials
5 which contain these compounds. Exemplary nitrogen sources include ammonia gas or ammonia salts, such as NH_4Cl or $(\text{NH}_4)_2\text{SO}_4$, NH_4OH , nitrates, urea, amino acids or complex nitrogen sources like corn steep liquor, soy bean flour, soy bean protein, yeast extract, meat extract and others.

Inorganic salt compounds which may be included in the media include the
10 chloride-, phosphorous- or sulfate- salts of calcium, magnesium, sodium, cobalt, molybdenum, potassium, manganese, zinc, copper and iron. Chelating compounds can be added to the medium to keep the metal ions in solution. Particularly useful chelating compounds include dihydroxyphenols, like catechol or protocatechuate, or organic acids, such as citric acid. It is typical for the media to also contain other growth factors, such as
15 vitamins or growth promoters, examples of which include biotin, riboflavin, thiamin, folic acid, nicotinic acid, pantothenate and pyridoxin. Growth factors and salts frequently originate from complex media components such as yeast extract, molasses, corn steep liquor and others. The exact composition of the media compounds depends strongly on the immediate experiment and is individually decided for each specific case. Information
20 about media optimization is available in the textbook "Applied Microbiol. Physiology, A Practical Approach (*eds.* P.M. Rhodes, P.F. Stanbury, IRL Press (1997) pp. 53-73, ISBN 0 19 963577 3). It is also possible to select growth media from commercial suppliers, like standard 1 (Merck) or BHI (grain heart infusion, DIFCO) or others.

All medium components are sterilized, either by heat (20 minutes at 1.5 bar and
25 121°C) or by sterile filtration. The components can either be sterilized together or, if necessary, separately. All media components can be present at the beginning of growth, or they can optionally be added continuously or batchwise.

Culture conditions are defined separately for each experiment. The temperature should be in a range between 15°C and 45°C. The temperature can be kept constant or can
30 be altered during the experiment. The pH of the medium should be in the range of 5 to 8.5, preferably around 7.0, and can be maintained by the addition of buffers to the media. An exemplary buffer for this purpose is a potassium phosphate buffer. Synthetic buffers such as MOPS, HEPES, ACES and others can alternatively or simultaneously be used. It is also possible to maintain a constant culture pH through the addition of NaOH or
35 NH_4OH during growth. If complex medium components such as yeast extract are utilized, the necessity for additional buffers may be reduced, due to the fact that many complex

compounds have high buffer capacities. If a fermentor is utilized for culturing the micro-organisms, the pH can also be controlled using gaseous ammonia.

The incubation time is usually in a range from several hours to several days. This time is selected in order to permit the maximal amount of product to accumulate in the
5 broth. The disclosed growth experiments can be carried out in a variety of vessels, such as microtiter plates, glass tubes, glass flasks or glass or metal fermentors of different sizes. For screening a large number of clones, the microorganisms should be cultured in microtiter plates, glass tubes or shake flasks, either with or without baffles. Preferably
10 100 ml shake flasks are used, filled with 10% (by volume) of the required growth medium. The flasks should be shaken on a rotary shaker (amplitude 25 mm) using a speed-range of 100 – 300 rpm. Evaporation losses can be diminished by the maintenance of a humid atmosphere; alternatively, a mathematical correction for evaporation losses should be performed.

If genetically modified clones are tested, an unmodified control clone or a control
15 clone containing the basic plasmid without any insert should also be tested. The medium is inoculated to an OD₆₀₀ of 0.5 – 1.5 using cells grown on agar plates, such as CM plates (10 g/l glucose, 2.5 g/l NaCl, 2 g/l urea, 10 g/l polypeptone, 5 g/l yeast extract, 5 g/l meat extract, 22 g/l NaCl, 2 g/l urea, 10 g/l polypeptone, 5 g/l yeast extract, 5 g/l meat extract, 22 g/l agar, pH 6.8 with 2M NaOH) that had been incubated at 30°C. Inoculation of the
20 media is accomplished by either introduction of a saline suspension of *C. glutamicum* cells from CM plates or addition of a liquid preculture of this bacterium.

Example 8 – *In vitro* Analysis of the Function of Mutant Proteins

The determination of activities and kinetic parameters of enzymes is well
25 established in the art. Experiments to determine the activity of any given altered enzyme must be tailored to the specific activity of the wild-type enzyme, which is well within the ability of one of ordinary skill in the art. Overviews about enzymes in general, as well as specific details concerning structure, kinetics, principles, methods, applications and examples for the determination of many enzyme activities may be
30 found, for example, in the following references: Dixon, M., and Webb, E.C., (1979) Enzymes. Longmans: London; Fersht, (1985) Enzyme Structure and Mechanism. Freeman: New York; Walsh, (1979) Enzymatic Reaction Mechanisms. Freeman: San Francisco; Price, N.C., Stevens, L. (1982) Fundamentals of Enzymology. Oxford Univ. Press: Oxford; Boyer, P.D., ed. (1983) The Enzymes, 3rd ed. Academic Press: New
35 York; Bisswanger, H., (1994) Enzymkinetik, 2nd ed. VCH: Weinheim (ISBN 3527300325); Bergmeyer, H.U., Bergmeyer, J., Graßl, M., eds. (1983-1986) Methods of Enzymatic Analysis, 3rd ed., vol. I-XII, Verlag Chemie: Weinheim; and Ullmann's

Encyclopedia of Industrial Chemistry (1987) vol. A9, "Enzymes". VCH: Weinheim, p. 352-363.

The activity of proteins which bind to DNA can be measured by several well-established methods, such as DNA band-shift assays (also called gel retardation assays).

5 The effect of such proteins on the expression of other molecules can be measured using reporter gene assays (such as that described in Kolmar, H. *et al.* (1995) *EMBO J.* 14: 3895-3904 and references cited therein). Reporter gene test systems are well known and established for applications in both pro- and eukaryotic cells, using enzymes such as beta-galactosidase, green fluorescent protein, and several others.

10 The determination of activity of membrane-transport proteins can be performed according to techniques such as those described in Gennis, R.B. (1989) "Pores, Channels and Transporters", in *Biomembranes, Molecular Structure and Function*, Springer: Heidelberg, p. 85-137; 199-234; and 270-322.

15 **Example 9: Analysis of Impact of Mutant Protein on the Production of the Desired Product**

The effect of the genetic modification in *C. glutamicum* on production of a desired compound (such as an amino acid) can be assessed by growing the modified microorganism under suitable conditions (such as those described above) and analyzing
20 the medium and/or the cellular component for increased production of the desired product (*i.e.*, an amino acid). Such analysis techniques are well known to one of ordinary skill in the art, and include spectroscopy, thin layer chromatography, staining methods of various kinds, enzymatic and microbiological methods, and analytical chromatography such as high performance liquid chromatography (see, for example,
25 Ullman, Encyclopedia of Industrial Chemistry, vol. A2, p. 89-90 and p. 443-613, VCH: Weinheim (1985); Fallon, A. *et al.*, (1987) "Applications of HPLC in Biochemistry" in: Laboratory Techniques in Biochemistry and Molecular Biology, vol. 17; Rehm *et al.* (1993) *Biotechnology*, vol. 3, Chapter III: "Product recovery and purification", page 469-714, VCH: Weinheim; Belter, P.A. *et al.* (1988) *Bioseparations: downstream*
30 processing for biotechnology, John Wiley and Sons; Kennedy, J.F. and Cabral, J.M.S. (1992) *Recovery processes for biological materials*, John Wiley and Sons; Shaeiwitz, J.A. and Henry, J.D. (1988) *Biochemical separations*, in: *Ullmann's Encyclopedia of Industrial Chemistry*, vol. B3, Chapter 11, page 1-27, VCH: Weinheim; and Dechow, F.J. (1989) *Separation and purification techniques in biotechnology*, Noyes
35 Publications.)

In addition to the measurement of the final product of fermentation, it is also possible to analyze other components of the metabolic pathways utilized for the

production of the desired compound, such as intermediates and side-products, to determine the overall yield, production, and/or efficiency of production of the compound. Analysis methods include measurements of nutrient levels in the medium (e.g., sugars, hydrocarbons, nitrogen sources, phosphate, and other ions), measurements of biomass composition and growth, analysis of the production of common metabolites of biosynthetic pathways, and measurement of gasses produced during fermentation. Standard methods for these measurements are outlined in Applied Microbial Physiology, A Practical Approach, P.M. Rhodes and P.F. Stanbury, eds., IRL Press, p. 103-129; 131-163; and 165-192 (ISBN: 0199635773) and references cited therein.

Example 10: Purification of the Desired Product from *C. glutamicum* Culture

Recovery of the desired product from the *C. glutamicum* cells or supernatant of the above-described culture can be performed by various methods well known in the art. If the desired product is not secreted from the cells, the cells can be harvested from the culture by low-speed centrifugation, the cells can be lysed by standard techniques, such as mechanical force or sonication. The cellular debris is removed by centrifugation, and the supernatant fraction containing the soluble proteins is retained for further purification of the desired compound. If the product is secreted from the *C. glutamicum* cells, then the cells are removed from the culture by low-speed centrifugation, and the supernate fraction is retained for further purification.

The supernatant fraction from either purification method is subjected to chromatography with a suitable resin, in which the desired molecule is either retained on a chromatography resin while many of the impurities in the sample are not, or where the impurities are retained by the resin while the sample is not. Such chromatography steps may be repeated as necessary, using the same or different chromatography resins. One of ordinary skill in the art would be well-versed in the selection of appropriate chromatography resins and in their most efficacious application for a particular molecule to be purified. The purified product may be concentrated by filtration or ultrafiltration, and stored at a temperature at which the stability of the product is maximized.

There are a wide array of purification methods known to the art and the preceding method of purification is not meant to be limiting. Such purification techniques are described, for example, in Bailey, J.E. & Ollis, D.F. Biochemical Engineering Fundamentals, McGraw-Hill: New York (1986).

The identity and purity of the isolated compounds may be assessed by techniques standard in the art. These include high-performance liquid chromatography (HPLC), spectroscopic methods, staining methods, thin layer chromatography, NIRS, enzymatic assay, or microbiologically. Such analysis methods are reviewed in: Patek *et al.* (1994)

- Appl. Environ. Microbiol.* 60: 133-140; Malakhova *et al.* (1996) *Biotechnologiya* 11: 27-32; and Schmidt *et al.* (1998) *Bioprocess Engineer.* 19: 67-70. Ulmann's Encyclopedia of Industrial Chemistry, (1996) vol. A27, VCH: Weinheim, p. 89-90, p. 521-540, p. 540-547, p. 559-566, 575-581 and p. 581-587; Michal, G. (1999) *Biochemical Pathways: An Atlas of Biochemistry and Molecular Biology*, John Wiley and Sons; Fallon, A. *et al.* (1987) *Applications of HPLC in Biochemistry in: Laboratory Techniques in Biochemistry and Molecular Biology*, vol. 17.

EXAMPLE 11: Cloning of a *Corynebacterium glutamicum* Gene Involved in Lincomycin Resistance Using a Reporter Gene Approach

A. Identification of the Gene Encoding the LMRB Protein

Plasmid pSL130 was constructed by ligation of the aceB promoter region (paceB) of *C. glutamicum* (Kim, H.J. *et al.* (1997) *J. Microbiol. Biotechnol.* 7: 287-292) into the polylinker of the lac operon fusion vector pRS415, which lacks a promoter (Simon, R.W. *et al.* (1987) *Gene* 53: 85-96). Plasmid pSL145 was constructed by ligating the resulting paceB-lac region into the *E. coli* cloning vector pACYC184. *E. coli* DH5aF' was transformed with pSL145 and the resulting strain was used as a host for screening of a genomic *C. glutamicum* library (in pSL109).

Transformants were screened by growth on agar medium containing 5-bromo-4-chloro-3-indolyl-beta-D-galactopyranoside (X-Gal). A white colony, containing DNA influencing lacZ expression, was selected for further analysis. This clone was found to contain a 4 kB fragment from the gene library. Subclones were constructed in pSL109 and a subclone which retained the white phenotype on X-Gal plates was identified. This subclone was found to contain a 2.6 kB BamH1-XhoI fragment (plasmid pSL149-5). The fragment was sequenced and identified as a membrane protein-encoding gene (LMRB gene).

The 1442 nucleotides of the coding sequence of the LMRB gene encode a polypeptide of 481 amino acid residues with a high percentage of hydrophobic amino acids. A Genbank search determined that the LMRB protein is 40% identical to the protein product of the lmrB gene from *Bacillus subtilis* (Genbank Accession AL009126, TREMBL Accession P94422), as determined using a CLUSTAL W analysis (using standard parameters).

The LMRN protein contains a sequence pattern: 158-A-P-A-L-G-P-T-L-S-G-167 (SEQ ID NO:301), which resembles the known multi-drug-resistance-protein consensus motif G-X-X-X-G-P-X-X-G-G (SEQ ID NO:302) (Paulsen, I.T., and Skurray, R.A.

(1993) *Gene* 124: 1-11). Therefore, the LMRB protein was classified as a drug resistance protein.

B. In vivo Analysis of lmrB Function

5 The *lmrB* gene was overexpressed in *C. glutamicum* ASO19E12 (Kim, H.J. *et al.* (1997) *J. Microbiol. Biotechnol.* 7: 287-292) using the plasmid pSL149-5, described above.

Disruption of the LMRB gene was accomplished by use of the vector pSL18-
lmrB. This vector was constructed as follows: an internal fragment of the LMRB gene
10 was amplified by PCR under standard conditions using primers 5'-
CTCCAGGATTGCTCCGAAGG-3' (SEQ ID NO:303) and 5'-
CACAGTGGTTGACCACTGGC-3' (SEQ ID NO:304). The resulting PCR product
was treated with T7 DNA polymerase and T7 polynucleotide kinase, and was cloned
into the *Sma*I site of plasmid pSL18 (Kim, Y.H. and H.-S. Lee (1996) *J. Microbiol.*
15 *Biotechnol.* 6: 315-320). The disruption of the LMRB gene in *C. glutamicum*
ASO19E12 was performed by conjugation, as previously described (Schwarzer and
Puhler (1991) *Bio/Technology* 9:84-87).

C. glutamicum cells transformed with pSL149-5 displayed similar resistances as
untransformed cells against erythromycin, penicillin G, tetracycline, chloramphenicol,
20 spectinomycin, nalidixic acid, gentamycin, streptomycin, ethidium bromide, carbonyl
cyanide m-chlorophenylhydrazone (CCCP), and sodium dodecyl sulfate. Significant
differences were observed, however, in the resistance of transformed and untransformed
cells to lincomycin.

LMRB-overexpressing *C. glutamicum* cells were found to be able to grow in the
25 presence of 20 µg/ml lincomycin. In contrast, cells which do not overexpress LMRB (or
cells carrying a LMRB disruption) were not able to grow on agar media containing 5
µg/ml lincomycin. This effect was clearly visible in liquid culture. LMRB
overexpression led to a 9-fold increased resistance (compared to wild-type) against
lincomycin and LMRB disruption resulted in a decreased resistance (28% of wild-type)
30 to this antibiotic.

Example 12: Analysis of the Gene Sequences of the Invention

The comparison of sequences and determination of percent homology between
two sequences are art-known techniques, and can be accomplished using a mathematical
35 algorithm, such as the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci.*
USA 87:2264-68, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci.* USA
90:5873-77. Such an algorithm is incorporated into the NBLAST and XBLAST

programs (version 2.0) of Altschul, *et al.* (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to SRT nucleic acid molecules of the invention. BLAST protein searches can be performed with the
5 XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to SRT protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped
10 BLAST programs, one of ordinary skill in the art will know how to optimize the parameters of the program (*e.g.*, XBLAST and NBLAST) for the specific sequence being analyzed.

Another example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Meyers and Miller ((1988) *Comput. Appl. Biosci.* 4: 11-17). Such an algorithm is incorporated into the ALIGN program (version 2.0) which is
15 part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used. Additional algorithms for sequence analysis are known in the art, and include ADVANCE and ADAM. described in Torelli and Robotti (1994) *Comput. Appl. Biosci.* 10:3-5; and FASTA, described in
20 Pearson and Lipman (1988) *P.N.A.S.* 85:2444-8.

The percent homology between two amino acid sequences can also be accomplished using the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blosum 62 matrix or a PAM250 matrix, and a gap weight of 12, 10, 8, 6, or 4 and a length weight of 2, 3, or 4. The percent homology
25 between two nucleic acid sequences can be accomplished using the GAP program in the GCG software package, using standard parameters, such as a gap weight of 50 and a length weight of 3.

A comparative analysis of the gene sequences of the invention with those present in Genbank has been performed using techniques known in the art (see, *e.g.*, Bexevanis and Ouellette, eds. (1998) *Bioinformatics: A Practical Guide to the Analysis of Genes and Proteins.* John Wiley and Sons: New York). The gene sequences of the invention were compared to genes present in Genbank in a three-step process. In a first step, a
30 BLASTN analysis (*e.g.*, a local alignment analysis) was performed for each of the sequences of the invention against the nucleotide sequences present in Genbank, and the top 500 hits were retained for further analysis. A subsequent FASTA search (*e.g.*, a combined local and global alignment analysis, in which limited regions of the sequences are aligned) was performed on these 500 hits. Each gene sequence of the invention was
35

09603208-062300

subsequently globally aligned to each of the top three FASTA hits, using the GAP program in the GCG software package (using standard parameters). In order to obtain correct results, the length of the sequences extracted from Genbank were adjusted to the length of the query sequences by methods well-known in the art. The results of this analysis are set forth in Table 4. The resulting data is identical to that which would have been obtained had a GAP (global) analysis alone been performed on each of the genes of the invention in comparison with each of the references in Genbank, but required significantly reduced computational time as compared to such a database-wide GAP (global) analysis. Sequences of the invention for which no alignments above the cutoff values were obtained are indicated on Table 4 by the absence of alignment information. It will further be understood by one of ordinary skill in the art that the GAP alignment homology percentages set forth in Table 4 under the heading "% homology (GAP)" are listed in the European numerical format, wherein a ',' represents a decimal point. For example, a value of "40,345" in this column represents "40.345%".

Example 13: Construction and Operation of DNA Microarrays

The sequences of the invention may additionally be used in the construction and application of DNA microarrays (the design, methodology, and uses of DNA arrays are well known in the art, and are described, for example, in Schena, M. *et al.* (1995) *Science* 270: 467-470; Wodicka, L. *et al.* (1997) *Nature Biotechnology* 15: 1359-1367; DeSaizieu, A. *et al.* (1998) *Nature Biotechnology* 16: 45-48; and DeRisi, J.L. *et al.* (1997) *Science* 278: 680-686).

DNA microarrays are solid or flexible supports consisting of nitrocellulose, nylon, glass, silicone, or other materials. Nucleic acid molecules may be attached to the surface in an ordered manner. After appropriate labeling, other nucleic acids or nucleic acid mixtures can be hybridized to the immobilized nucleic acid molecules, and the label may be used to monitor and measure the individual signal intensities of the hybridized molecules at defined regions. This methodology allows the simultaneous quantification of the relative or absolute amount of all or selected nucleic acids in the applied nucleic acid sample or mixture. DNA microarrays, therefore, permit an analysis of the expression of multiple (as many as 6800 or more) nucleic acids in parallel (see, *e.g.*, Schena, M. (1996) *BioEssays* 18(5): 427-431).

The sequences of the invention may be used to design oligonucleotide primers which are able to amplify defined regions of one or more *C. glutamicum* genes by a nucleic acid amplification reaction such as the polymerase chain reaction. The choice and design of the 5' or 3' oligonucleotide primers or of appropriate linkers allows the covalent attachment of the resulting PCR products to the surface of a support medium

described above (and also described, for example, Schena, M. *et al.* (1995) *Science* 270: 467-470).

Nucleic acid microarrays may also be constructed by *in situ* oligonucleotide synthesis as described by Wodicka, L. *et al.* (1997) *Nature Biotechnology* 15: 1359-1367. By photolithographic methods, precisely defined regions of the matrix are exposed to light. Protective groups which are photolabile are thereby activated and undergo nucleotide addition, whereas regions that are masked from light do not undergo any modification. Subsequent cycles of protection and light activation permit the synthesis of different oligonucleotides at defined positions. Small, defined regions of the genes of the invention may be synthesized on microarrays by solid phase oligonucleotide synthesis.

The nucleic acid molecules of the invention present in a sample or mixture of nucleotides may be hybridized to the microarrays. These nucleic acid molecules can be labeled according to standard methods. In brief, nucleic acid molecules (*e.g.*, mRNA molecules or DNA molecules) are labeled by the incorporation of isotopically or fluorescently labeled nucleotides, *e.g.*, during reverse transcription or DNA synthesis. Hybridization of labeled nucleic acids to microarrays is described (*e.g.*, in Schena, M. *et al.* (1995) *supra*; Wodicka, L. *et al.* (1997), *supra*; and DeSaizieu A. *et al.* (1998), *supra*). The detection and quantification of the hybridized molecule are tailored to the specific incorporated label. Radioactive labels can be detected, for example, as described in Schena, M. *et al.* (1995) *supra*) and fluorescent labels may be detected, for example, by the method of Shalon *et al.* (1996) *Genome Research* 6: 639-645).

The application of the sequences of the invention to DNA microarray technology, as described above, permits comparative analyses of different strains of *C. glutamicum* or other Corynebacteria. For example, studies of inter-strain variations based on individual transcript profiles and the identification of genes that are important for specific and/or desired strain properties such as pathogenicity, productivity and stress tolerance are facilitated by nucleic acid array methodologies. Also, comparisons of the profile of expression of genes of the invention during the course of a fermentation reaction are possible using nucleic acid array technology.

Example 14: Analysis of the Dynamics of Cellular Protein Populations (Proteomics)

The genes, compositions, and methods of the invention may be applied to study the interactions and dynamics of populations of proteins, termed 'proteomics'. Protein populations of interest include, but are not limited to, the total protein population of *C. glutamicum* (*e.g.*, in comparison with the protein populations of other organisms), those

proteins which are active under specific environmental or metabolic conditions (e.g., during fermentation, at high or low temperature, or at high or low pH), or those proteins which are active during specific phases of growth and development.

Protein populations can be analyzed by various well-known techniques, such as gel electrophoresis. Cellular proteins may be obtained, for example, by lysis or extraction, and may be separated from one another using a variety of electrophoretic techniques. Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) separates proteins largely on the basis of their molecular weight. Isoelectric focusing polyacrylamide gel electrophoresis (IEF-PAGE) separates proteins by their isoelectric point (which reflects not only the amino acid sequence but also posttranslational modifications of the protein). Another, more preferred method of protein analysis is the consecutive combination of both IEF-PAGE and SDS-PAGE, known as 2-D-gel electrophoresis (described, for example, in Hermann *et al.* (1998) *Electrophoresis* 19: 3217-3221; Fountoulakis *et al.* (1998) *Electrophoresis* 19: 1193-1202; Langen *et al.* (1997) *Electrophoresis* 18: 1184-1192; Antelmann *et al.* (1997) *Electrophoresis* 18: 1451-1463). Other separation techniques may also be utilized for protein separation, such as capillary gel electrophoresis; such techniques are well known in the art.

Proteins separated by these methodologies can be visualized by standard techniques, such as by staining or labeling. Suitable stains are known in the art, and include Coomassie Brilliant Blue, silver stain, or fluorescent dyes such as Sypro Ruby (Molecular Probes). The inclusion of radioactively labeled amino acids or other protein precursors (e.g., ^{35}S -methionine, ^{35}S -cysteine, ^{14}C -labelled amino acids, ^{15}N -amino acids, $^{15}\text{NO}_3$ or $^{15}\text{NH}_4^+$ or ^{13}C -labelled amino acids) in the medium of *C. glutamicum* permits the labeling of proteins from these cells prior to their separation. Similarly, fluorescent labels may be employed. These labeled proteins can be extracted, isolated and separated according to the previously described techniques.

Proteins visualized by these techniques can be further analyzed by measuring the amount of dye or label used. The amount of a given protein can be determined quantitatively using, for example, optical methods and can be compared to the amount of other proteins in the same gel or in other gels. Comparisons of proteins on gels can be made, for example, by optical comparison, by spectroscopy, by image scanning and analysis of gels, or through the use of photographic films and screens. Such techniques are well-known in the art.

To determine the identity of any given protein, direct sequencing or other standard techniques may be employed. For example, N- and/or C-terminal amino acid sequencing (such as Edman degradation) may be used, as may mass spectrometry (in particular MALDI or ESI techniques (see, e.g., Langen *et al.* (1997) *Electrophoresis* 18:

09603208-062300

The information obtained by these methods can be used to compare patterns of protein presence, activity, or modification between different samples from various biological conditions (*e.g.*, different organisms, time points of fermentation, media conditions, or different biotopes, among others). Data obtained from such experiments alone, or in combination with other techniques, can be used for various applications, such as to compare the behavior of various organisms in a given (*e.g.*, metabolic) situation, to increase the productivity of strains which produce fine chemicals or to increase the efficiency of the production of fine chemicals.

Those of ordinary skill in the art will recognize, or will be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

[illegible]



TABLE 1: Genes Included in the Application

<u>Nucleic Acid SEQ ID NO</u>	<u>Amino Acid SEQ ID NO</u>	<u>Identification Code</u>	<u>Contig.</u>	<u>NT Start</u>	<u>NT Stop</u>	<u>Function</u>
1	2	RXA01524	GR00424	29041	30483	Lincomycin RESISTANCE PROTEIN
3	4	RXA00497	GR00124	52	348	10 KD CHAPERONIN
5	6	RXN00493	VV0086	14389	16002	60 KD CHAPERONIN
7	8	F RXA00498	GR00124	363	1601	60 KD CHAPERONIN
9	10	RXA01217	GR00353	802	203	GENERAL STRESS PROTEIN CTC
11	12	RXA00605	GR00159	7412	5865	CATALASE (EC 1.11.1.6)
13	14	RXA00404	GR00089	2909	594	CARBON STARVATION PROTEIN A
15	16	RXN03119	VV0098	86877	87008	SUPEROXIDE DISMUTASE [MN] (EC 1.15.1.1)
17	18	RXN03120	VV0098	87351	87476	SUPEROXIDE DISMUTASE [MN] (EC 1.15.1.1)
19	20	RXN00575	VV0323	14716	15252	PHOSPHINOTHRICIN-RESISTANCE PROTEIN
21	22	F RXA00575	GR00156	2130	1648	PHOSPHINOTHRICIN-RESISTANCE PROTEIN

Chaperones

<u>Nucleic Acid SEQ ID NO</u>	<u>Amino Acid SEQ ID NO</u>	<u>Identification Code</u>	<u>Contig.</u>	<u>NT Start</u>	<u>NT Stop</u>	<u>Function</u>
23	24	RXN01345	VV0123	4883	3432	Molecular chaperon (HSP70/Dnak family)
25	26	F RXA01345	GR00391	1172	6	Molecular chaperones (HSP70/Dnak family)
27	28	RXA02541	GR00726	13657	12473	DNAJ PROTEIN
29	30	RXA02542	GR00726	14518	13865	GRPE PROTEIN
31	32	RXN02543	VV0057	22031	20178	DNAK PROTEIN
33	34	F RXA02543	GR00726	16375	14522	DNAK PROTEIN
35	36	RXN02280	VV0152	1849	26	TRAP1
37	38	F RXA02282	GR00659	1145	1480	Molecular chaperone, HSP90 family
39	40	RXA00886	GR00242	12396	13541	DNAJ PROTEIN
41	42	RXS00568	VV0251	2928	1582	TRIGGER FACTOR
43	44	RXN03038	VV0017	42941	43666	PS1 PROTEIN VORLAUFER
45	46	RXN03039	VV0018	2	631	PS1 PROTEIN VORLAUFER
47	48	RXN03040	VV0018	761	1069	PS1 PROTEIN VORLAUFER
49	50	RXN03051	VV0022	2832	3566	PS1 PROTEIN VORLAUFER
51	52	RXN03054	VV0026	1906	3486	PS1 PROTEIN VORLAUFER
53	54	RXN02949	VV0025	31243	31575	PREPROTEIN TRANSLOKASE SECE UNTEREINHEIT
55	56	RXN02462	VV0124	11932	13749	PREPROTEIN TRANSLOKASE SECA UNTEREINHEIT
57	58	RXN01559	VV0171	7795	5954	PROTEIN-EXPORT MEMBRANE PROTEIN SEC2
59	60	RXN00046	VV0119	5363	6058	Signal Erkennung particle GTPase
61	62	RXN01863	VV0206	1172	24	/O/C Thioedoxin-ähnliche oxidoreductase
63	64	RXN00833	VV0180	8039	8533	THIOL PEROXIDASE (EC 1.11.1.-)

<u>Nucleic Acid SEQ ID NO</u>	<u>Amino Acid SEQ ID NO</u>	<u>Identification Code</u>	<u>Contig.</u>	<u>NT Start</u>	<u>NT Stop</u>	<u>Function</u>
65	66	RXN01676	VV0179	12059	11304	THIOL-DISULFIDE AUSTAUSCH PROTEIN DSBD
67	68	RXN00380	VV0223	836	216	THIOL-DISULFIDE AUSTAUSCH PROTEIN TLPA
69	70	RXN00937	VV0079	42335	42706	THIOREDOXIN
71	72	RXN02325	VV0047	5527	6393	THIOREDOXIN
73	74	RXN01837	VV0320	7103	7879	PEPTIDYL-PROLYL CIS-TRANS ISOMERASE (EC 5.2.1.8)
75	76	RXN01926	VV0284	1	741	PEPTID KETTE RELEASE FACTOR 3
77	78	RXN02002	VV0111	141	518	PEPTID KETTE RELEASE FACTOR 3
79	80	RXN02736	VV0074	13600	14556	PUTATIVES OXPPCYCLE PROTEIN OPCA
81	82	RXS03217				SMALL COLD-SHOCK PROTEIN
83	84	F RXA01917	GR00549	3465	3665	SMALL COLD-SHOCK PROTEIN

Proteins involved in stress responses

<u>Nucleic Acid SEQ ID NO</u>	<u>Amino Acid SEQ ID NO</u>	<u>Identification Code</u>	<u>Contig.</u>	<u>NT Start</u>	<u>NT Stop</u>	<u>Function</u>
85	86	RXA02184	GR00641	19628	19248	COLD SHOCK-LIKE PROTEIN CSPC
87	88	RXA00810	GR00218	792	992	SMALL COLD-SHOCK PROTEIN
89	90	RXA01674	GR00467	1878	2771	PROBABLE HYDROGEN PEROXIDE-INDUCIBLE GENES ACTIVATOR
91	92	RXA02431	GR00708	2	1192	damage-inducible protein P
93	94	RXA02446	GR00709	11640	11206	OSMOTICALLY INDUCIBLE PROTEIN C
95	96	RXA02861	GR10006	551	1633	probable metallothionein u0308aa - Mycobacterium leprae
97	98	RXA00981	GR00276	3388	4017	GTP PYROPHOSPHOKINASE (EC 2.7.6.5)
99	100	RXN00786	VV0321	1680	706	LYTB PROTEIN
101	102	RXS01027	VV0143	5761	6768	DIADENOSINE 5',5'''-P1,P4-TETRAPHOSPHATE HYDROLASE (EC 3.6.1.17)
103	104	RXS01528	VV0050	17276	16749	DIADENOSINE 5',5'''-P1,P4-TETRAPHOSPHATE HYDROLASE (EC 3.6.1.17)
105	106	RXS01716	VV0319	3259	2774	EXOPOLYPHOSPHATASE (EC 3.6.1.11)
107	108	RXS01835	VV0143	10575	10045	GUANOSINE-3',5'-BIS(DIPHOSPHATE) 3'-PYROPHOSPHOHYDROLASE (EC 3.1.7.2)
109	110	RXS02497	VV0007	15609	16535	EXOPOLYPHOSPHATASE (EC 3.6.1.11)
111	112	RXS02972	VV0319	2763	2353	EXOPOLYPHOSPHATASE (EC 3.6.1.11)

Resistance and tolerance

<u>Nucleic Acid SEQ ID NO</u>	<u>Amino Acid SEQ ID NO</u>	<u>Identification Code</u>	<u>Contig.</u>	<u>NT Start</u>	<u>NT Stop</u>	<u>Function</u>
113	114	RXA02159	GR00640	6231	6743	ARGININE HYDROXIMATE RESISTANCE PROTEIN
115	116	RXA02201	GR00646	5837	6199	ARSENATE REDUCTASE
117	118	RXA00599	GR00159	1843	1457	ARSENICAL-RESISTANCE PROTEIN ACR3
119	120	RXA00600	GR00159	2940	1843	ARSENICAL-RESISTANCE PROTEIN ACR3
121	122	RXA02200	GR00646	4651	5760	ARSENICAL-RESISTANCE PROTEIN ACR3
123	124	RXA02202	GR00646	6278	6916	ARSENICAL-RESISTANCE PROTEIN ACR3
125	126	RXA02205	GR00646	9871	8993	BACITRACIN RESISTANCE PROTEIN (PUTATIVE UNDECAPRENOL KINASE) (EC 2.7.1.66)
127	128	RXA00900	GR00245	4052	3201	BICYCLOWYCIN RESISTANCE PROTEIN

<u>Nucleic Acid SEQ ID NO</u>	<u>Amino Acid SEQ ID NO</u>	<u>Identification Code</u>	<u>Contig.</u>	<u>NT Start</u>	<u>NT Stop</u>	<u>Function</u>
129	130	RXN00901	VV0140	8581	8168	BICYCLOMYCIN RESISTANCE PROTEIN
131	132	F RXA00901	GR00245	4357	3980	BICYCLOMYCIN RESISTANCE PROTEIN
133	134	RXA00289	GR00046	3263	4438	CHLORAMPHENICOL RESISTANCE PROTEIN
135	136	RXN01984	VV0056	1515	1811	CHLORAMPHENICOL RESISTANCE PROTEIN
137	138	F RXA01984	GR00574	282	4	CHLORAMPHENICOL RESISTANCE PROTEIN
139	140	RXA00109	GR00015	1176	565	COPPER RESISTANCE PROTEIN C PRECURSOR
141	142	RXA00109	GR00015	1176	565	COPPER RESISTANCE PROTEIN C PRECURSOR
143	144	RXA00996	GR00283	1763	1023	DAUNORUBICIN RESISTANCE ATP-BINDING PROTEIN DRRA
145	146	RXN00829	VV0180	7950	5611	DAUNORUBICIN RESISTANCE PROTEIN
147	148	F RXA00829	GR00224	2	256	DAUNORUBICIN RESISTANCE PROTEIN
149	150	F RXA00834	GR00225	463	2025	DAUNORUBICIN RESISTANCE PROTEIN
151	152	RXA00995	GR00283	1023	283	DAUNORUBICIN RESISTANCE TRANSMEMBRANE PROTEIN
153	154	RXN00803	VV0009	53858	52629	METHYLENOMYCIN A RESISTANCE PROTEIN
155	156	F RXA00803	GR00214	4560	5162	METHYLENOMYCIN A RESISTANCE PROTEIN
157	158	RXA01407	GR00410	3918	3028	METHYLENOMYCIN A RESISTANCE PROTEIN
159	160	RXA01408	GR00410	4384	4184	METHYLENOMYCIN A RESISTANCE PROTEIN
161	162	RXN01922	VV0020	2031	3182	METHYLENOMYCIN A RESISTANCE PROTEIN
163	164	F RXA01922	GR00552	3	1109	METHYLENOMYCIN A RESISTANCE PROTEIN
165	166	RXA02060	GR00626	1	339	MYCINAMICIN-RESISTANCE PROTEIN MYRA
167	168	RXN01936	VV0127	40116	41387	MACROLIDE-EFFLUX PROTEIN
169	170	F RXA01936	GR00555	9796	8975	NICKEL RESISTANCE PROTEIN
171	172	F RXA01937	GR00555	10246	9821	NICKEL RESISTANCE PROTEIN
173	174	RXN01010	VV0209	3776	4894	QUINOLONE RESISTANCE NORA PROTEIN
175	176	F RXA01010	GR00288	774	4	QUINOLONE RESISTANCE NORA PROTEIN
177	178	RXN03142	VV0136	5754	4612	QUINOLONE RESISTANCE NORA PROTEIN
179	180	F RXA01150	GR00323	3807	2917	QUINOLONE RESISTANCE NORA PROTEIN
181	182	RXN02964	VV0102	7931	6714	QUINOLONE RESISTANCE NORA PROTEIN
183	184	F RXA02116	GR00636	911	6	QUINOLONE RESISTANCE NORA PROTEIN
185	186	RXA00858	GR00233	1680	2147	TELLURIUM RESISTANCE PROTEIN TERC
187	188	RXA02305	GR00663	2921	2070	DAUNOMYCIN C-14 HYDROXYLASE
189	190	RXA00084	GR00013	2367	1543	VIBRIOBACTIN UTILIZATION PROTEIN VIUB
191	192	RXA00843	GR00228	3236	3580	ARSENATE REDUCTASE
193	194	RXA01052	GR00296	3398	3706	MERCURIC REDUCTASE (EC 1.16.1.1)
195	196	RXA01053	GR00296	3772	4191	MERCURIC REDUCTASE (EC 1.16.1.1)
197	198	RXA01054	GR00296	4229	4717	MERCURIC REDUCTASE (EC 1.16.1.1)
199	200	RXN03123	VV0106	808	1245	HEAVY METAL TOLERANCE PROTEIN PRECURSOR
201	202	F RXA00993	GR00282	641	6	HEAVY METAL TOLERANCE PROTEIN PRECURSOR
203	204	RXA01051	GR00296	3298	2690	VANZ PROTEIN , teicoplanin resistance protein
205	206	RXN01873	VV0248	2054	819	Hypothetical Drug Resistance Protein
207	208	F RXA01873	GR00535	855	1946	Hypothetical Drug Resistance Protein
209	210	RXN00034	VV0020	16933	18381	MULTIDRUG RESISTANCE PROTEIN B
211	212	F RXA02273	GR00655	8058	9002	Hypothetical Drug Resistance Protein
213	214	RXN03075	VV0042	2491	3216	Hypothetical Drug Transporter
215	216	F RXA02907	GR10044	1395	2120	Hypothetical Drug Transporter
217	218	RXA00479	GR00119	16290	14101	Hypothetical Drug Transporter
219	220	RXN03124	VV0108	4	963	Hypothetical Drug Transporter
221	222	F RXA01180	GR00336	4	765	Hypothetical Drug Transporter

Table 1, Page 3

<u>Nucleic Acid SEQ ID NO</u>	<u>Amino Acid SEQ ID NO</u>	<u>Identification Code</u>	<u>Contig.</u>	<u>NT Start</u>	<u>NT Stop</u>	<u>Function</u>
223	224	RXA02586	GR00741	10296	10027	Hypothetical Drug Transporter
225	226	RXA02587	GR00741	12343	10253	Hypothetical Drug Transporter
227	228	RXN03042	VV0018	2440	1835	Hypothetical Drug Transporter
229	230	F RXA02893	GR10035	1841	1236	Hypothetical Drug Transporter
231	232	RXA01616	GR00450	1684	203	MULTIDRUG EFFLUX PROTEIN QACB
233	234	RXA01666	GR00463	2307	3683	MULTIDRUG RESISTANCE PROTEIN
235	236	RXA00062	GR00009	13252	11855	MULTIDRUG RESISTANCE PROTEIN B
237	238	RXA00215	GR00032	13834	15294	MULTIDRUG RESISTANCE PROTEIN B
239	240	RXN03064	VV0038	4892	6223	MULTIDRUG RESISTANCE PROTEIN B
241	242	F RXA00565	GR00151	4892	5884	MULTIDRUG RESISTANCE PROTEIN B
243	244	F RXA02878	GR10016	1837	1481	MULTIDRUG RESISTANCE PROTEIN B
245	246	RXA00648	GR00169	2713	1304	MULTIDRUG RESISTANCE PROTEIN B
247	248	RXN01320	VV0082	13146	11500	MULTIDRUG RESISTANCE PROTEIN B
249	250	F RXA01314	GR00382	744	4	MULTIDRUG RESISTANCE PROTEIN B
251	252	F RXA01320	GR00383	1979	1200	MULTIDRUG RESISTANCE PROTEIN B
253	254	RXN02926	VV0082	11497	9866	MULTIDRUG RESISTANCE PROTEIN B
255	256	F RXA01319	GR00383	1197	4	MULTIDRUG RESISTANCE PROTEIN B
257	258	RXA01578	GR00439	1423	29	MULTIDRUG RESISTANCE PROTEIN B
259	260	RXA02087	GR00629	7076	5730	MULTIDRUG RESISTANCE PROTEIN B
261	262	RXA02088	GR00629	8294	7080	MULTIDRUG RESISTANCE PROTEIN B
263	264	RXA00764	GR00204	3284	2169	BMRU PROTEIN <i>Bacillus subtilis</i> bmrU, multidrug efflux transporter
265	266	RXN03125	VV0108	972	1142	Hypothetical Drug Transporter
267	268	RXN01553	VV0135	25201	26520	Hypothetical Drug Permease
269	270	RXN00535	VV0219	5155	5871	Hypothetical Drug Resistance Protein
271	272	RXN00453	VV0076	1173	3521	Hypothetical Drug Transporter
273	274	RXN00932	VV0171	13120	13593	Hypothetical Drug Transporter
275	276	RXN03022	VV0002	65	511	MULTIDRUG RESISTANCE PROTEIN B
277	278	RXN03151	VV0163	489	4	MYCINAMICIN-RESISTANCE PROTEIN MYRA
279	280	RXN02832	VV0358	547	5	LYSOSTAPHIN IMMUNITY FACTOR
281	282	RXN00165	VV0232	3275	1860	MULTIDRUG RESISTANCE-LIKE ATP-BINDING PROTEIN MDL
283	284	RXN01190	VV0169	8992	10338	MULTIDRUG RESISTANCE-LIKE ATP-BINDING PROTEIN MDL
285	286	RXN01102	VV0059	6128	4884	QUINOLONE RESISTANCE NORA PROTEIN
287	288	RXN00788	VV0321	3424	3648	CHLORAMPHENICOL RESISTANCE PROTEIN
289	290	RXN02119	VV0102	11242	9602	AZ01A-RESISTANCE ATP-BINDING PROTEIN
291	292	RXN01605	VV0137	7124	5610	DAUNORUBICIN RESISTANCE TRANSMEMBRANE PROTEIN
293	294	RXN01091	VV0326	567	4	MAZG PROTEIN
295	296	RXS02979	VV0149	2150	2383	MERCURIC TRANSPORT PROTEIN PERIPLASMIC COMPONENT PRECURSOR
297	298	RXS02987	VV0234	527	294	MERCURIC TRANSPORT PROTEIN PERIPLASMIC COMPONENT PRECURSOR
299	300	RXS03095	VV0057	4056	4424	CADMIUM EFFLUX SYSTEM ACCESSORY PROTEIN HOMOLOG



00/02/90

80209/60
old 'S' n 8580f

Attorney Docket No.: BGI-124CP

GenBank™ Accession No.	Gene Name	Gene Function	Reference
A09073	ppg	Phosphoenol pyruvate carboxylase	Bachmann, B. et al. "DNA fragment coding for phosphoenolpyruvate carboxylase, recombinant DNA carrying said fragment, strains carrying the recombinant DNA and method for producing L-aminino acids using said strains," Patent: EP 0358940-A 3 03/21/90
A45579, A45581, A45583, A45585 A45587		Threonine dehydratase	Moeckel, B. et al. "Production of L-isoleucine by means of recombinant micro-organisms with deregulated threonine dehydratase," Patent: WO 9519442-A 5 07/20/95
AB003132	murC; ftsQ; ftsZ		Kobayashi, M. et al. "Cloning, sequencing, and characterization of the ftsZ gene from coryneform bacteria," <i>Biochem. Biophys. Res. Commun.</i> , 236(2):383-388 (1997)
AB015023	murC; ftsQ		Wachi, M. et al. "A murC gene from Coryneform bacteria," <i>Appl. Microbiol. Biotechnol.</i> , 51(2):223-228 (1999)
AB018530	disR		Kimura, E. et al. "Molecular cloning of a novel gene, disR, which rescues the detergent sensitivity of a mutant derived from <i>Brevibacterium lactofermentum</i> ," <i>Biosci. Biotechnol. Biochem.</i> , 60(10):1565-1570 (1996)
AB018531	disR1; disR2		
AB020624	murI	D-glutamate racemase	
AB023377	tkl	transketolase	
AB024708	gltB; gltD	Glutamine 2-oxoglutarate aminotransferase large and small subunits	
AB025424	acn	aconitase	
AB027714	rep	Replication protein	
AB027715	rep; aad	Replication protein; aminoglycoside adenylyltransferase	
AF005242	argC	N-acetylglutamate-5-semialdehyde dehydrogenase	
AF005635	glnA	Glutamine synthetase	
AF030405	hisF	cyclase	
AF030520	argG	Argininosuccinate synthetase	
AF031518	argF	Ornithine carbamoyltransferase	
AF036932	aroD	3-dehydroquinate dehydratase	

GenBank™ Accession No.	Gene Name	Gene Function	Reference
AF038548	pyc	Pyruvate carboxylase	
AF038651	dcIAE; apt; rel	Dipeptide-binding protein; adenine phosphoribosyltransferase; GTP pyrophosphokinase	Wehmeier, L. et al. "The role of the <i>Corynebacterium glutamicum</i> rel gene in (p)ppGpp metabolism," <i>Microbiology</i> , 144:1853-1862 (1998)
AF041436	argR	Arginine repressor	
AF045998	impA	Inositol monophosphate phosphatase	
AF048764	argH	Argininosuccinate lyase	
AF049897	argC; argI; argB; argD; argF; argR; argG; argH	N-acetylglutanylphosphate reductase; ornithine acetyltransferase; N-acetylglutamate kinase; acetylornithine transaminase; ornithine carbamoyltransferase; arginine repressor; argininosuccinate synthase; argininosuccinate lyase	
AF050109	inhA	Enoyl-acyl carrier protein reductase	
AF050166	hisG	ATP phosphoribosyltransferase	
AF051846	hisA	Phosphoribosylformimino-5-amino-1-phosphoribosyl-4-imidazolecarboxamide isomerase	
AF052652	metA	Homoserine O-acetyltransferase	Park, S. et al. "Isolation and analysis of metA, a methionine biosynthetic gene encoding homoserine acetyltransferase in <i>Corynebacterium glutamicum</i> ," <i>Mol. Cells</i> , 8(3):286-294 (1998)
AF053071	aroB	Dehydroquinate synthetase	
AF060558	hisH	Glutamine amidotransferase	
AF086704	hisE	Phosphoribosyl-ATP-pyrophosphohydrolase	
AF114233	aroA	5-enolpyruvylshikimate 3-phosphate synthase	
AF116184	pand	L-aspartate-alpha-decarboxylase precursor	Dusch, N. et al. "Expression of the <i>Corynebacterium glutamicum</i> pand gene encoding L-aspartate-alpha-decarboxylase leads to pantothenate overproduction in <i>Escherichia coli</i> ," <i>Appl. Environ. Microbiol.</i> , 65(4):1530-1539 (1999)

GenBank™ Accession No.	Gene Name	Gene Function	Reference
AF124518	aroD; aroE	3-dehydroquinase; shikimate dehydrogenase	
AF124600	aroC; aroK; aroB; pepQ	Chorismate synthase; shikimate kinase; 3-dehydroquinase synthase; putative cytoplasmic peptidase	
AF145897	inhA		
AF145898	inhA		
AJ001436	ectP	Transport of ectoine, glycine betaine, proline	Peter, H. et al. "Corynebacterium glutamicum is equipped with four secondary carriers for compatible solutes: Identification, sequencing, and characterization of the proline/ectoine uptake system, ProP, and the ectoine/proline/glycine betaine carrier, EctP," <i>J. Bacteriol.</i> , 180(22):6005-6012 (1998)
AJ004934	dapD	Tetrahydrodipicolinate succinylase (incomplete)	Wehrmann, A. et al. "Different modes of diaminopimelate synthesis and their role in cell wall integrity: A study with Corynebacterium glutamicum," <i>J. Bacteriol.</i> , 180(12):3159-3165 (1998)
AJ007732	ppe; secG; amt; ocd; soxA	Phosphoenolpyruvate-carboxylase; ?, high affinity ammonium uptake protein; putative ornithine-cyclodecarboxylase; sarcosine oxidase	
AJ010319	ftsY; glnB, glnD; srp; amtP	Involved in cell division; PII protein; uridylyltransferase (uridylyl-removing enzyme); signal recognition particle; low affinity ammonium uptake protein	Jakoby, M. et al. "Nitrogen regulation in Corynebacterium glutamicum; Isolation of genes involved in biochemical characterization of corresponding proteins," <i>FEMS Microbiol.</i> , 173(2):303-310 (1999)
AJ132968	cat	Chloramphenicol acetyl transferase	
AJ224946	mgo	L-malate: quinone oxidoreductase	Molenaar, D. et al. "Biochemical and genetic characterization of the membrane-associated malate dehydrogenase (acceptor) from Corynebacterium glutamicum," <i>Eur. J. Biochem.</i> , 254(2):395-403 (1998)
AJ238250	ndh	NADH dehydrogenase	
AJ238703	porA	Porin	Lichtinger, T. et al. "Biochemical and biophysical characterization of the cell wall porin of Corynebacterium glutamicum: The channel is formed by a low molecular mass polypeptide," <i>Biochemistry</i> , 37(43):15024-15032 (1998)
DI7429		Transposable element IS31831	Vertes, A.A. et al. "Isolation and characterization of IS31831, a transposable element from Corynebacterium glutamicum," <i>Mol. Microbiol.</i> , 11(4):739-746 (1994)

GenBank™ Accession No.	Gene Name	Gene Function	Reference
D84102	odhA	2-oxoglutarate dehydrogenase	Usuda, Y. et al. "Molecular cloning of the <i>Corynebacterium glutamicum</i> (Brevibacterium lactofermentum A12036) <i>odhA</i> gene encoding a novel type of 2-oxoglutarate dehydrogenase," <i>Microbiology</i> , 142:3347-3354 (1996)
E01358	hdh; hk	Homoserine dehydrogenase; homoserine kinase	Katsumata, R. et al. "Production of L-threonine and L-isoleucine," Patent: JP 1987232392-A 1 10/12/87
E01359		Upstream of the start codon of homoserine kinase gene	Katsumata, R. et al. "Production of L-threonine and L-isoleucine," Patent: JP 1987232392-A 2 10/12/87
E01375		Tryptophan operon	
E01376	trpL; trpE	Leader peptide; anthranilate synthase	Matsui, K. et al. "Tryptophan operon, peptide and protein coded thereby, utilization of tryptophan operon gene expression and production of tryptophan," Patent: JP 1987244382-A 1 10/24/87
E01377		Promoter and operator regions of tryptophan operon	Matsui, K. et al. "Tryptophan operon, peptide and protein coded thereby, utilization of tryptophan operon gene expression and production of tryptophan," Patent: JP 1987244382-A 1 10/24/87
E03937		Biotin-synthase	Hatakeyama, K. et al. "DNA fragment containing gene capable of coding biotin synthetase and its utilization," Patent: JP 1992278088-A 1 10/02/92
E04040		Diamino pelargonic acid aminotransferase	Kohama, K. et al. "Gene coding diaminopelargonic acid aminotransferase and deshiobiotin synthetase and its utilization," Patent: JP 1992330284-A 1 11/18/92
E04041		Deshiobiotinsynthetase	Kohama, K. et al. "Gene coding diaminopelargonic acid aminotransferase and deshiobiotin synthetase and its utilization," Patent: JP 1992330284-A 1 11/18/92
E04307		Flavum aspartase	Kurusu, Y. et al. "Gene DNA coding aspartase and utilization thereof," Patent: JP 1993030977-A 1 02/09/93
E04376		Isocitric acid lyase	Katsumata, R. et al. "Gene manifestation controlling DNA," Patent: JP 1993056782-A 3 03/09/93
E04377		Isocitric acid lyase N-terminal fragment	Katsumata, R. et al. "Gene manifestation controlling DNA," Patent: JP 1993056782-A 3 03/09/93
E04484		Prephenate dehydratase	Sotouchi, N. et al. "Production of L-phenylalanine by fermentation," Patent: JP 1993076352-A 2 03/30/93
E05108		Aspartokinase	Fugono, N. et al. "Gene DNA coding Aspartokinase and its use," Patent: JP 1993184366-A 1 07/27/93
E05112		Dihydro-dipicolinate synthetase	Hatakeyama, K. et al. "Gene DNA coding dihydrodipicolinic acid synthetase and its use," Patent: JP 1993184371-A 1 07/27/93

GenBank™ Accession No.	Gene Name	Gene Function	Reference
E05776		Diaminopimelic acid dehydrogenase	Kobayashi, M. et al. "Gene DNA coding Diaminopimelic acid dehydrogenase and its use," Patent: JP 1993284970-A 1 11/02/93
E05779		Threonine synthase	Kohama, K. et al. "Gene DNA coding threonine synthase and its use," Patent: JP 1993284972-A 1 11/02/93
E06110		Prephenate dehydratase	Kikuchi, T. et al. "Production of L-phenylalanine by fermentation method," Patent: JP 1993344881-A 1 12/27/93
E06111		Mutated Prephenate dehydratase	Kikuchi, T. et al. "Production of L-phenylalanine by fermentation method," Patent: JP 1993344881-A 1 12/27/93
E06146		Acetohydroxy acid synthetase	Inui, M. et al. "Gene capable of coding Acetohydroxy acid synthetase and its use," Patent: JP 1993344893-A 1 12/27/93
E06825		Aspartokinase	Sugimoto, M. et al. "Mutant aspartokinase gene," patent: JP 1994062866-A 1 03/08/94
E06826		Mutated aspartokinase alpha subunit	Sugimoto, M. et al. "Mutant aspartokinase gene," patent: JP 1994062866-A 1 03/08/94
E06827		Mutated aspartokinase alpha subunit	Sugimoto, M. et al. "Mutant aspartokinase gene," patent: JP 1994062866-A 1 03/08/94
E07701	secY		Honno, N. et al. "Gene DNA participating in integration of membraneous protein to membrane," Patent: JP 1994169780-A 1 06/21/94
E08177		Aspartokinase	Sato, Y. et al. "Genetic DNA capable of coding Aspartokinase released from feedback inhibition and its utilization," Patent: JP 1994261766-A 1 09/20/94
E08178, E08179, E08180, E08181, E08182		Feedback inhibition-released Aspartokinase	Sato, Y. et al. "Genetic DNA capable of coding Aspartokinase released from feedback inhibition and its utilization," Patent: JP 1994261766-A 1 09/20/94
E08232		Acetohydroxy-acid isomeroeductase	Inui, M. et al. "Gene DNA coding acetohydroxy acid isomeroeductase," Patent: JP 1994277067-A 1 10/04/94
E08234	secE		Asai, Y. et al. "Gene DNA coding for translocation machinery of protein," Patent: JP 1994277073-A 1 10/04/94
E08643		FT aminotransferase and desthiobiotin synthetase promoter region	Hatakeyama, K. et al. "DNA fragment having promoter function in coryneform bacterium," Patent: JP 1995031476-A 1 02/03/95
E08646		Biotin synthetase	Hatakeyama, K. et al. "DNA fragment having promoter function in coryneform bacterium," Patent: JP 1995031476-A 1 02/03/95

GenBank™ Accession No.	Gene Name	Gene Function	Reference
E08649		Aspartase	Kohama, K. et al "DNA fragment having promoter function in coryneform bacterium," Patent: JP 1995031478-A 1 02/03/95
E08900		Dihydrodipicolinate reductase	Madori, M. et al. "DNA fragment containing gene coding Dihydrodipicolinate acid reductase and utilization thereof," Patent: JP 1995075578-A 1 03/20/95
E08901		Diaminopimelic acid decarboxylase	Madori, M. et al. "DNA fragment containing gene coding Diaminopimelic acid decarboxylase and utilization thereof," Patent: JP 1995075579-A 1 03/20/95
E12594		Serine hydroxymethyltransferase	Hatakeyama, K. et al. "Production of L-tryptophan," Patent: JP 1997028391-A 1 02/04/97
E12760, E12759, E12758		transposase	Moriya, M. et al. "Amplification of gene using artificial transposon," Patent: JP 1997070291-A 03/18/97
E12764		Arginyl-tRNA synthetase; diaminopimelic acid decarboxylase	Moriya, M. et al. "Amplification of gene using artificial transposon," Patent: JP 1997070291-A 03/18/97
E12767		Dihydrodipicolinic acid synthetase	Moriya, M. et al. "Amplification of gene using artificial transposon," Patent: JP 1997070291-A 03/18/97
E12770		aspartokinase	Moriya, M. et al. "Amplification of gene using artificial transposon," Patent: JP 1997070291-A 03/18/97
E12773		Dihydrodipicolinic acid reductase	Moriya, M. et al. "Amplification of gene using artificial transposon," Patent: JP 1997070291-A 03/18/97
E13655		Glucose-6-phosphate dehydrogenase	Hatakeyama, K. et al. "Glucose-6-phosphate dehydrogenase and DNA capable of coding the same," Patent: JP 1997224661-A 1 09/02/97
L01508	IlvA	Threonine dehydratase	Moeckel, B. et al. "Functional and structural analysis of the threonine dehydratase of <i>Corynebacterium glutamicum</i> ," <i>J. Bacteriol.</i> , 174:8065-8072 (1992)
L07603	EC 4.2.1.15	3-deoxy-D-arabinoheptulosonate-7-phosphate synthase	Chen, C. et al. "The cloning and nucleotide sequence of <i>Corynebacterium glutamicum</i> 3-deoxy-D-arabinoheptulosonate-7-phosphate synthase gene," <i>FEMS Microbiol. Lett.</i> , 107:223-230 (1993)
L09232	IlvB; ilvN; ilvC	Acetohydroxy acid synthase large subunit; Acetohydroxy acid synthase small subunit; Acetohydroxy acid isomerase	Keilhauer, C. et al. "Isolencine synthesis in <i>Corynebacterium glutamicum</i> : molecular analysis of the ilvB-ilvN-ilvC operon," <i>J. Bacteriol.</i> , 175(17):5595-5603 (1993)

GenBank™ Accession No.	Gene Name	Gene Function	Reference
L18874	ptsM	Phosphoenolpyruvate sugar phosphotransferase	Fouet, A et al. "Bacillus subtilis sucrose-specific enzyme II of the phosphotransferase system: expression in <i>Escherichia coli</i> and homology to enzymes II from enteric bacteria," <i>PNAS USA</i> , 84(24):8773-8777 (1987); Lee, J.K. et al. "Nucleotide sequence of the gene encoding the <i>Corynebacterium glutamicum</i> mannose enzyme II and analyses of the deduced protein sequence," <i>FEMS Microbiol. Lett.</i> , 119(1-2):137-145 (1994)
L27123	aceB	Malate synthase	Lee, H-S. et al. "Molecular characterization of aceB, a gene encoding malate synthase in <i>Corynebacterium glutamicum</i> ," <i>J. Microbiol. Biotechnol.</i> , 4(4):256-263 (1994)
L27126		Pyruvate kinase	Jetten, M. S. et al. "Structural and functional analysis of pyruvate kinase from <i>Corynebacterium glutamicum</i> ," <i>Appl. Environ. Microbiol.</i> , 60(7):2501-2507 (1994)
L28760	aceA	Isocitrate lyase	
L35906	dxr	Diphtheria toxin repressor	Oguiza, J.A. et al. "Molecular cloning, DNA sequence analysis, and characterization of the <i>Corynebacterium diphtheriae</i> dxr from <i>Brevibacterium lactofermentum</i> ," <i>J. Bacteriol.</i> , 177(2):465-467 (1995)
M13774		Prephenate dehydratase	Folletite, M.T. et al. "Molecular cloning and nucleotide sequence of the <i>Corynebacterium glutamicum</i> pheA gene," <i>J. Bacteriol.</i> , 167:695-702 (1986)
M16175	5S rRNA		Park, Y-H. et al. "Phylogenetic analysis of the coryneform bacteria by 5S rRNA sequences," <i>J. Bacteriol.</i> , 169:1801-1806 (1987)
M16663	tpuE	Anthraniolate synthase, 5' end	Sano, K. et al. "Structure and function of the trp operon control regions of <i>Brevibacterium lactofermentum</i> , a glutamic-acid-producing bacterium," <i>Gene</i> , 52:191-200 (1987)
M16664	tpuA	Tryptophan synthase, 3' end	Sano, K. et al. "Structure and function of the trp operon control regions of <i>Brevibacterium lactofermentum</i> , a glutamic-acid-producing bacterium," <i>Gene</i> , 52:191-200 (1987)
M25819		Phosphoenolpyruvate carboxylase	O'Regan, M. et al. "Cloning and nucleotide sequence of the Phosphoenolpyruvate carboxylase-coding gene of <i>Corynebacterium glutamicum</i> ATCC13032," <i>Gene</i> , 77(2):237-251 (1989)
M85106		23S rRNA gene insertion sequence	Roller, C. et al. "Gram-positive bacteria with a high DNA G+C content are characterized by a common insertion within their 23S rRNA genes," <i>J. Gen. Microbiol.</i> , 138:1167-1175 (1992)

GenBank™ Accession No.	Gene Name	Gene Function	Reference
M85107, M85108		23S rRNA gene insertion sequence	Roller, C. et al. "Gram-positive bacteria with a high DNA G+C content are characterized by a common insertion within their 23S rRNA genes," <i>J. Gen. Microbiol.</i> , 138:1167-1175 (1992)
M89931	aecD; brnQ; yhbW	Beta C-S lyase; branched-chain amino acid uptake carrier; hypothetical protein yhbW	Rosso, I. et al. "The <i>Corynebacterium glutamicum</i> aecD gene encodes a C-S lyase with alpha, beta-elimination activity that degrades aminocetylcyserine," <i>J. Bacteriol.</i> , 174(9):2968-2977 (1992); Tauch, A. et al. "Isolucine uptake in <i>Corynebacterium glutamicum</i> ATCC 13032 is directed by the brnQ gene product," <i>Arch. Microbiol.</i> , 169(4):303-312 (1998)
S59299	trp	Leader gene (promoter)	Herry, D.M. et al. "Cloning of the trp gene cluster from a tryptophan-hyperproducing strain of <i>Corynebacterium glutamicum</i> : identification of a mutation in the trp leader sequence," <i>Appl. Environ. Microbiol.</i> , 59(3):791-799 (1993)
U11545	trpD	Anthranilate phosphoribosyltransferase	O'Gara, J.P. and Dunican, L.K. (1994) Complete nucleotide sequence of the <i>Corynebacterium glutamicum</i> ATCC 21850 trpD gene." Thesis, Microbiology Department, University College Galway, Ireland.
U13922	cgIIIM; cgIIIR; cglIIIR	Putative type II 5-cytosine methyltransferase; putative type II restriction endonuclease; putative type I or type III restriction endonuclease	Schafer, A. et al. "Cloning and characterization of a DNA region encoding a stress-sensitive restriction system from <i>Corynebacterium glutamicum</i> ATCC 13032 and analysis of its role in intergeneric conjugation with <i>Escherichia coli</i> ," <i>J. Bacteriol.</i> , 176(23):7309-7319 (1994); Schafer, A. et al. "The <i>Corynebacterium glutamicum</i> cgIIIM gene encoding a 5-cytosine in an MrcBC-deficient <i>Escherichia coli</i> strain," <i>Gene</i> , 203(2):95-101 (1997)
U14965	recA		
U31224	ppx		Ankri, S. et al. "Mutations in the <i>Corynebacterium glutamicum</i> proline biosynthetic pathway: A natural bypass of the proA step," <i>J. Bacteriol.</i> , 178(15):4412-4419 (1996)
U31225	proC	L-proline: NADP+ 5-oxidoreductase	Ankri, S. et al. "Mutations in the <i>Corynebacterium glutamicum</i> proline biosynthetic pathway: A natural bypass of the proA step," <i>J. Bacteriol.</i> , 178(15):4412-4419 (1996)
U31230	obg; proB; unkdh	?; gamma glutamyl kinase; similar to D-isomer specific 2-hydroxyacid dehydrogenases	Ankri, S. et al. "Mutations in the <i>Corynebacterium glutamicum</i> proline biosynthetic pathway: A natural bypass of the proA step," <i>J. Bacteriol.</i> , 178(15):4412-4419 (1996)

GenBank™ Accession No.	Gene Name	Gene Function	Reference
U31281	bioB	Biotin synthase	Serebriiskii, I.G., "Two new members of the bio B superfamily: Cloning, sequencing and expression of bio B genes of <i>Methylobacillus flagellatum</i> and <i>Corynebacterium glutamicum</i> ," <i>Gene</i> , 175:15-22 (1996)
U35023	thrR; accBC	Thiosulfate sulfurtransferase; acyl CoA carboxylase	Jager, W. et al. "A <i>Corynebacterium glutamicum</i> gene encoding a two-domain protein similar to biotin carboxylases and biotin-carboxyl-carrier proteins," <i>Arch. Microbiol.</i> , 166(2):76-82 (1996)
U43535	cmr	Multidrug resistance protein	Jager, W. et al. "A <i>Corynebacterium glutamicum</i> gene conferring multidrug resistance in the heterologous host <i>Escherichia coli</i> ," <i>J. Bacteriol.</i> , 179(7):2449-2451 (1997)
U43536	clpB	Heat shock ATP-binding protein	
U53587	aphA-3	3'5'-aminoglycoside phosphotransferase	
U89648		<i>Corynebacterium glutamicum</i> unidentified sequence involved in histidine biosynthesis, partial sequence	
X04960	trpA; trpB; trpC; trpD; trpE; trpG; trpL	Tryptophan operon	Matsui, K. et al. "Complete nucleotide and deduced amino acid sequences of the <i>Brevibacterium lactofermentum</i> tryptophan operon," <i>Nucleic Acids Res.</i> , 14(24):10113-10114 (1986)
X07563	lys A	DAP decarboxylase (meso-diaminopimelate decarboxylase, EC 4.1.1.20)	Yeh, P. et al. "Nucleic sequence of the <i>lysA</i> gene of <i>Corynebacterium glutamicum</i> and possible mechanisms for modulation of its expression," <i>Mol. Gen. Genet.</i> , 212(1):112-119 (1988)
X14234	EC 4.1.1.31	Phosphoenolpyruvate carboxylase	Eikmanns, B.J. et al. "The Phosphoenolpyruvate carboxylase gene of <i>Corynebacterium glutamicum</i> : Molecular cloning, nucleotide sequence, and expression," <i>Mol. Gen. Genet.</i> , 218(2):330-339 (1989); Lepiniec, L. et al. "Sorghum Phosphoenolpyruvate carboxylase gene family: structure, function and molecular evolution," <i>Plant. Mol. Biol.</i> , 21 (3):487-502 (1993)
X17313	fda	Fructose-bisphosphate aldolase	Von der Osten, C.H. et al. "Molecular cloning, nucleotide sequence and fine-structural analysis of the <i>Corynebacterium glutamicum</i> <i>fda</i> gene: structural comparison of <i>C. glutamicum</i> fructose-1, 6-biphosphate aldolase to class I and class II aldolases," <i>Mol. Microbiol.</i>
X53993	dapA	L-2, 3-dihydrodipicolinate synthetase (EC 4.2.1.52)	Bonnassie, S. et al. "Nucleic sequence of the <i>dapA</i> gene from <i>Corynebacterium glutamicum</i> ," <i>Nucleic Acids Res.</i> , 18(21):6421 (1990)

GenBank™ Accession No.	Gene Name	Gene Function	Reference
X54223		AttB-related site	Cianciotto, N. et al. "DNA sequence homology between att B-related sites of <i>Corynebacterium diphtheriae</i> , <i>Corynebacterium ulcerans</i> , <i>Corynebacterium glutamicum</i> , and the attP site of <i>lambdacorynephage</i> ," <i>FEMS. Microbiol. Lett.</i> , 66:299-302 (1990)
X54740	argS, lysA	Arginyl-tRNA synthetase; Diaminopimelate decarboxylase	Marcel, T. et al. "Nucleotide sequence and organization of the upstream region of the <i>Corynebacterium glutamicum</i> lysA gene," <i>Mol. Microbiol.</i> , 4(11):1819-1830 (1990)
X55994	trpL, trpE	Putative leader peptide; anthranilate synthase component I	Heery, D.M. et al. "Nucleotide sequence of the <i>Corynebacterium glutamicum</i> trpE gene," <i>Nucleic Acids Res.</i> , 18(23):7138 (1990)
X56037	thrC	Threonine synthase	Han, K.S. et al. "The molecular structure of the <i>Corynebacterium glutamicum</i> threonine synthase gene," <i>Mol. Microbiol.</i> , 4(10):1693-1702 (1990)
X56075	attB-related site	Attachment site	Cianciotto, N. et al. "DNA sequence homology between att B-related sites of <i>Corynebacterium diphtheriae</i> , <i>Corynebacterium ulcerans</i> , <i>Corynebacterium glutamicum</i> , and the attP site of <i>lambdacorynephage</i> ," <i>FEMS. Microbiol. Lett.</i> , 66:299-302 (1990)
X57226	lysC-alpha; lysC-beta; asd	Aspartokinase-alpha subunit; Aspartokinase-beta subunit; aspartate beta semialdehyde dehydrogenase	Kalinowski, J. et al. "Genetic and biochemical analysis of the Aspartokinase from <i>Corynebacterium glutamicum</i> ," <i>Mol. Microbiol.</i> , 5(5):1197-1204 (1991); Kalinowski, J. et al. "Aspartokinase genes lysC alpha and lysC beta overlap and are adjacent to the aspartate beta-semialdehyde dehydrogenase gene asd in <i>Corynebacterium glutamicum</i> ," <i>Mol. Gen. Genet.</i> , 224(3):317-324 (1990)
X59403	gap;pgk; tpi	Glyceraldehyde-3-phosphate; phosphoglycerate kinase; triosephosphate isomerase	Eikmanns, B.J. "Identification, sequence analysis, and expression of a <i>Corynebacterium glutamicum</i> gene cluster encoding the three glycolytic enzymes glyceraldehyde-3-phosphate dehydrogenase, 3-phosphoglycerate kinase, and triosephosphate isomerases," <i>J. Bacteriol.</i> , 174(19):6076-6086 (1992)
X59404	gdh	Glutamate dehydrogenase	Bormann, E.R. et al. "Molecular analysis of the <i>Corynebacterium glutamicum</i> gdh gene encoding glutamate dehydrogenase," <i>Mol. Microbiol.</i> , 6(3):317-326 (1992)
X60312	lysI	L-lysine permease	Seep-Feldhaus, A.H. et al. "Molecular analysis of the <i>Corynebacterium glutamicum</i> lysI gene involved in lysine uptake," <i>Mol. Microbiol.</i> , 5(12):2995-3005 (1991)

GenBank™ Accession No.	Gene Name	Gene Function	Reference
X66078	copI	PSI protein	Joliff, G. et al. "Cloning and nucleotide sequence of the cspI gene encoding PSI, one of the two major secreted proteins of <i>Corynebacterium glutamicum</i> : The deduced N-terminal region of PSI is similar to the <i>Mycobacterium</i> antigen 85 complex," <i>Mol. Microbiol.</i> , 6(16):2349-2362 (1992)
X66112	glt	Citrate synthase	Eikmanns, B.J. et al. "Cloning sequence, expression and transcriptional analysis of the <i>Corynebacterium glutamicum</i> gltA gene encoding citrate synthase," <i>Microbiol.</i> , 140:1817-1828 (1994)
X67737	dapB	Dihydropicolinate reductase	
X69103	csp2	Surface layer protein PS2	Peyret, J.L. et al. "Characterization of the cspB gene encoding PS2, an ordered surface-layer protein in <i>Corynebacterium glutamicum</i> ," <i>Mol. Microbiol.</i> , 9(1):97-109 (1993)
X69104		IS3 related insertion element	Bonamy, C. et al. "Identification of IS1206, a <i>Corynebacterium glutamicum</i> IS3-related insertion sequence and phylogenetic analysis," <i>Mol. Microbiol.</i> , 14(3):571-581 (1994)
X70959	leuA	Isopropylmalate synthase	Patek, M. et al. "Leucine synthesis in <i>Corynebacterium glutamicum</i> : enzyme activities, structure of leuA, and effect of leuA inactivation on lysine synthesis," <i>Appl. Environ. Microbiol.</i> , 60(1):133-140 (1994)
X71489	icd	Isocitrate dehydrogenase (NADP+)	Elkmanns, B.J. et al. "Cloning sequence analysis, expression, and inactivation of the <i>Corynebacterium glutamicum</i> icd gene encoding isocitrate dehydrogenase and biochemical characterization of the enzyme," <i>J. Bacteriol.</i> , 177(3):774-782 (1995)
X72855	GDHA	Glutamate dehydrogenase (NADP+)	
X75083, X70584	mitA	5-methyltryptophan resistance	Heery, D.M. et al. "A sequence from a tryptophan-hyperproducing strain of <i>Corynebacterium glutamicum</i> encoding resistance to 5-methyltryptophan," <i>Biochem. Biophys. Res. Commun.</i> , 201(3):1255-1262 (1994)
X75085	recA		Fitzpatrick, R. et al. "Construction and characterization of recA mutant strains of <i>Corynebacterium glutamicum</i> and <i>Brevibacterium lactofermentum</i> ," <i>Appl. Microbiol. Biotechnol.</i> , 42(4):575-580 (1994)
X75504	aceA; thiX	Partial Isocitrate lyase; ?	Reinscheid, D.J. et al. "Characterization of the isocitrate lyase gene from <i>Corynebacterium glutamicum</i> and biochemical analysis of the enzyme," <i>J. Bacteriol.</i> , 176(12):3474-3483 (1994)
X76875		ATPase beta-subunit	Ludwig, W. et al. "Phylogenetic relationships of bacteria based on comparative sequence analysis of elongation factor Tu and ATP-synthase beta-subunit genes," <i>Antonie Van Leeuwenhoek</i> , 64:285-305 (1993)

GenBank™ Accession No.	Gene Name	Gene Function	Reference
X77034	tuf	Elongation factor Tu	Ludwig, W. et al. "Phylogenetic relationships of bacteria based on comparative sequence analysis of elongation factor Tu and A TP-synthase beta-subunit genes," <i>Antonie Van Leeuwenhoek</i> , 64:285-305 (1993)
X77384	recA		Billman-Jacobe, H. "Nucleotide sequence of a recA gene from <i>Corynebacterium glutamicum</i> ," <i>DNA Seq.</i> , 4(6):403-404 (1994)
X78491	aceB	Malate synthase	Reinscheid, D.J. et al. "Malate synthase from <i>Corynebacterium glutamicum</i> pia-ack operon encoding phosphotransacetylase: sequence analysis," <i>Microbiology</i> , 140:3099-3108 (1994)
X80629	16S rDNA	16S ribosomal RNA	Railey, F.A. et al. "Phylogenetic analysis of the genera <i>Rhodococcus</i> and <i>Norcardia</i> and evidence for the evolutionary origin of the genus <i>Norcardia</i> from within the radiation of <i>Rhodococcus</i> species," <i>Microbiol.</i> , 141:523-528 (1995)
X81191	gluA; gluB; gluC; gluD	Glutamate uptake system	Kronmeyer, W. et al. "Structure of the gluABCD cluster encoding the glutamate uptake system of <i>Corynebacterium glutamicum</i> ," <i>J. Bacteriol.</i> , 177(5):1152-1158 (1995)
X81379	dapE	Succinylidiaininopimelate desuccinylase	Wehrmann, A. et al. "Analysis of different DNA fragments of <i>Corynebacterium glutamicum</i> complementing dapE of <i>Escherichia coli</i> ," <i>Microbiology</i> , 40:3349-56 (1994)
X82061	16S rDNA	16S ribosomal RNA	Ruimy, R. et al. "Phylogeny of the genus <i>Corynebacterium</i> deduced from analyses of small-subunit ribosomal DNA sequences," <i>Int. J. Syst. Bacteriol.</i> , 45(4):740-746 (1995)
X82928	asd; lysC	Aspartate-semialdehyde dehydrogenase; ?	Serebrijski, I. et al. "Multicopy suppression by asd gene and osmotic stress-dependent complementation by heterologous proA in proA mutants," <i>J. Bacteriol.</i> , 177(24):7255-7260 (1995)
X82929	proA	Gamma-glutamyl phosphate reductase	Serebrijski, I. et al. "Multicopy suppression by asd gene and osmotic stress-dependent complementation by heterologous proA in proA mutants," <i>J. Bacteriol.</i> , 177(24):7255-7260 (1995)
X84257	16S rDNA	16S ribosomal RNA	Pascual, C. et al. "Phylogenetic analysis of the genus <i>Corynebacterium</i> based on 16S rRNA gene sequences," <i>Int. J. Syst. Bacteriol.</i> , 45(4):724-728 (1995)
X85965	aroP; dapE	Aromatic amino acid permease; ?	Wehrmann, A. et al. "Functional analysis of sequences adjacent to dapE of <i>Corynebacterium glutamicum</i> reveals the presence of aroP, which encodes the aromatic amino acid transporter," <i>J. Bacteriol.</i> , 177(20):5991-5993 (1995)

GenBank™ Accession No.	Gene Name	Gene Function	Reference
X86157	argB; argC; argD; argF; argI	Acetylglutamate kinase; N-acetyl- γ -glutamyl-phosphate reductase; acetylornithine aminotransferase; ornithine carbamoyltransferase; glutamate N-acetyltransferase	Sakanyan, V. et al. "Genes and enzymes of the acetyl cycle of arginine biosynthesis in <i>Corynebacterium glutamicum</i> : enzyme evolution in the early steps of the arginine pathway," <i>Microbiology</i> , 142:99-108 (1996)
X89084	pta; ackA	Phosphate acetyltransferase; acetate kinase	Reinscheid, D.J. et al. "Cloning, sequence analysis, expression and inactivation of the <i>Corynebacterium glutamicum</i> pta-ack operon encoding phosphotransacetylase and acetate kinase," <i>Microbiology</i> , 145:503-513 (1999)
X89850	attB	Attachment site	Le Marrec, C. et al. "Genetic characterization of site-specific integration functions of phi AAU2 infecting "Arthrobacter aureus C70," <i>J. Bacteriol.</i> , 178(7):1996-2004 (1996)
X90356		Promoter fragment F1	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90357		Promoter fragment F2	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90358		Promoter fragment F10	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90359		Promoter fragment F13	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90360		Promoter fragment F22	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90361		Promoter fragment F34	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90362		Promoter fragment F37	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)

GenBank™ Accession No.	Gene Name	Gene Function	Reference
X90363		Promoter fragment F45	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90364		Promoter fragment F64	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90365		Promoter fragment F75	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90366		Promoter fragment PF101	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90367		Promoter fragment PF104	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90368		Promoter fragment PF109	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X93513	amt	Ammonium transport system	Siewe, R.M. et al. "Functional and genetic characterization of the (methyl) ammonium uptake carrier of <i>Corynebacterium glutamicum</i> ," <i>J. Biol. Chem.</i> , 271(10):5398-5403 (1996)
X93514	betP	Glycine betaine transport system	Peter, H. et al. "Isolation, characterization, and expression of the <i>Corynebacterium glutamicum</i> betP gene, encoding the transport system for the compatible solute glycine betaine," <i>J. Bacteriol.</i> , 178(17):5229-5234 (1996)
X95649	orf4		Patek, M. et al. "Identification and transcriptional analysis of the dapB-ORF2-dapA-ORF4 operon of <i>Corynebacterium glutamicum</i> , encoding two enzymes involved in L-lysine synthesis," <i>Biotechnol. Lett.</i> , 19:1113-1117 (1997)
X96471	lysE, lysG	Lysine exporter protein; Lysine export regulator protein	Vrjlic, M. et al. "A new type of transporter with a new type of cellular function: L-lysine export from <i>Corynebacterium glutamicum</i> ," <i>Mol. Microbiol.</i> , 22(5):815-826 (1996)

GenBank™ Accession No.	Gene Name	Gene Function	Reference
X96580	panB; panC; xylB	3-methyl-2-oxobutanoate hydroxymethyltransferase; pantoate-beta-alanine ligase; xylulokinase	Sahm, H. et al. "D-pantothenate synthesis in <i>Corynebacterium glutamicum</i> and use of panB and genes encoding L-valine synthesis for D-pantothenate overproduction," <i>Appl. Environ. Microbiol.</i> , 65(5):1973-1979 (1999)
X96962		Insertion sequence IS1207 and transposase	
X99289		Elongation factor P	Ramos, A. et al. "Cloning, sequencing and expression of the gene encoding elongation factor P in the amino-acid producer <i>Brevibacterium lactofermentum</i> (<i>Corynebacterium glutamicum</i> ATCC 13869)," <i>Gene</i> , 198:217-222 (1997)
Y00140	thrB	Homoserine kinase	Mateos, L.M. et al. "Nucleotide sequence of the homoserine kinase (thrB) gene of the <i>Brevibacterium lactofermentum</i> ," <i>Nucleic Acids Res.</i> , 15(9):3922 (1987)
Y00151	ddh	Meso-diaminopimelate D-dehydrogenase (EC 1.4.1.16)	Ishino, S. et al. "Nucleotide sequence of the meso-diaminopimelate D-dehydrogenase gene from <i>Corynebacterium glutamicum</i> ," <i>Nucleic Acids Res.</i> , 15(9):3917 (1987)
Y00476	thrA	Homoserine dehydrogenase	Mateos, L.M. et al. "Nucleotide sequence of the homoserine dehydrogenase (thrA) gene of the <i>Brevibacterium lactofermentum</i> ," <i>Nucleic Acids Res.</i> , 15(24):10598 (1987)
Y00546	hom; thrB	Homoserine dehydrogenase; homoserine kinase	Peoples, O.P. et al. "Nucleotide sequence and fine structural analysis of the <i>Corynebacterium glutamicum</i> hom-thrB operon," <i>Mol. Microbiol.</i> , 2(1):63-72 (1988)
Y08964	murC; ftsQ/divD; ftsZ	UPD-N-acetylmuramate-alanine ligase; division initiation protein or cell division protein; cell division protein	Hornubia, M.P. et al. "Identification, characterization, and chromosomal organization of the ftsZ gene from <i>Brevibacterium lactofermentum</i> ," <i>Mol. Gen. Genet.</i> , 259(1):97-104 (1998)
Y09163	putP	High affinity proline transport system	Peter, H. et al. "Isolation of the putP gene of <i>Corynebacterium glutamicum</i> proline and characterization of a low-affinity uptake system for compatible solutes," <i>Arch. Microbiol.</i> , 168(2):143-151 (1997)
Y09548	pyc	Pyruvate carboxylase	Peters-Wendisch, P.G. et al. "Pyruvate carboxylase from <i>Corynebacterium glutamicum</i> : characterization, expression and inactivation of the pyc gene," <i>Microbiology</i> , 144:915-927 (1998)
Y09578	leuB	3-isopropylmalate dehydrogenase	Patek, M. et al. "Analysis of the leuB gene from <i>Corynebacterium glutamicum</i> ," <i>Appl. Microbiol. Biotechnol.</i> , 50(1):42-47 (1998)
Y12472		Attachment site bacteriophage Phi-16	Moreau, S. et al. "Site-specific integration of coryneophage Phi-16: The construction of an integration vector," <i>Microbiol.</i> , 145:539-548 (1999)

GenBank™ Accession No.	Gene Name	Gene Function	Reference
Y12537	proP	Proline/ectoine uptake system protein	Peter, H. et al. "Corynebacterium glutamicum is equipped with four secondary carriers for compatible solutes: Identification, sequencing, and characterization of the proline/ectoine uptake system, ProP, and the ectoine/proline/glycine betaine carrier, EctP," <i>J. Bacteriol.</i> , 180(22):6005-6012 (1998)
Y13221	glnA	Glutamine synthetase I	Jakoby, M. et al. "Isolation of Corynebacterium glutamicum glnA gene encoding glutamine synthetase I," <i>FEMS Microbiol. Lett.</i> , 154(1):81-88 (1997)
Y16642	lpd	Dihydrolipoamide dehydrogenase	
Y18059		Attachment site Corynephage 304L	Moreau, S. et al. "Analysis of the integration functions of φ304L: An integrase module among corynephages," <i>Virology</i> , 255(1):150-159 (1999)
Z21501	argS, lysA	Arginyl-tRNA synthetase; diaminiopimelate decarboxylase (partial)	Oguiza, J.A. et al. "A gene encoding arginyl-tRNA synthetase is located in the upstream region of the lysA gene in Brevibacterium lactofermentum: Regulation of argS-lysA cluster expression by arginine," <i>J. Bacteriol.</i> , 175(22):7356-7362 (1993)
Z21502	dapA, dapB	Dihydrodipicolinate synthase; dihydrodipicolinate reductase	Pisabarro, A. et al. "A cluster of three genes (dapA, orf2, and dapB) of Brevibacterium lactofermentum encodes dihydrodipicolinate reductase, and a third polypeptide of unknown function," <i>J. Bacteriol.</i> , 175(9):2743-2749 (1993)
Z29563	thrC	Threonine synthase	Malumbres, M. et al. "Analysis and expression of the thrC gene of the encoded threonine synthase," <i>Appl. Environ. Microbiol.</i> , 60(7):2209-2219 (1994)
Z46753	16S rDNA	Gene for 16S ribosomal RNA	
Z49822	sigA	SigA sigma factor	Oguiza, J.A. et al. "Multiple sigma factor genes in Brevibacterium lactofermentum: Characterization of sigA and sigB," <i>J. Bacteriol.</i> , 178(2):550-553 (1996)
Z49823	galE, dtxR	Catalytic activity UDP-galactose 4-epimerase; diphtheria toxin regulatory protein	Oguiza, J.A. et al. "The galE gene encoding the UDP-galactose 4-epimerase of Brevibacterium lactofermentum is coupled transcriptionally to the dmdR gene," <i>Gene</i> , 177:103-107 (1996)
Z49824	orf1, sigB	?, SigB sigma factor	Oguiza, J.A. et al. "Multiple sigma factor genes in Brevibacterium lactofermentum: Characterization of sigA and sigB," <i>J. Bacteriol.</i> , 178(2):550-553 (1996)
Z66534		Transposase	Correia, A. et al. "Cloning and characterization of an IS-like element present in the genome of Brevibacterium lactofermentum ATCC 13869," <i>Gene</i> , 170(1):91-94 (1996)

A sequence for this gene was published in the indicated reference. However, the sequence obtained by the inventors of the present application is significantly longer than the published version. It is believed that the published version relied on an incorrect start codon, and thus represents only a fragment of the actual coding region.

TABLE 3: Corynebacterium and Brevibacterium Strains Which May be Used in the Practice of the Invention

Genus	Species	ATCC	FERM	NRRL	CECT	NCIMB	CBS	NCITC	DSMZ
Brevibacterium	ammoniagenes	21054							
Brevibacterium	ammoniagenes	19350							
Brevibacterium	ammoniagenes	19351							
Brevibacterium	ammoniagenes	19352							
Brevibacterium	ammoniagenes	19353							
Brevibacterium	ammoniagenes	19354							
Brevibacterium	ammoniagenes	19355							
Brevibacterium	ammoniagenes	19356							
Brevibacterium	ammoniagenes	21055							
Brevibacterium	ammoniagenes	21077							
Brevibacterium	ammoniagenes	21553							
Brevibacterium	ammoniagenes	21580							
Brevibacterium	ammoniagenes	39101							
Brevibacterium	butanicum	21196							
Brevibacterium	divaricatum	21792	P928						
Brevibacterium	flavum	21474							
Brevibacterium	flavum	21129							
Brevibacterium	flavum	21518							
Brevibacterium	flavum			B11474					
Brevibacterium	flavum			B11472					
Brevibacterium	flavum	21127							
Brevibacterium	flavum	21128							
Brevibacterium	flavum	21427							
Brevibacterium	flavum	21475							
Brevibacterium	flavum	21517							
Brevibacterium	flavum	21528							
Brevibacterium	flavum	21529							

Brevibacterium	flavum			B11477						
Brevibacterium	flavum			B11478						
Brevibacterium	flavum	21127								
Brevibacterium	flavum			B11474						
Brevibacterium	healii	15527								
Brevibacterium	ketoglutamicum	21004								
Brevibacterium	ketoglutamicum	21089								
Brevibacterium	ketosoreductum	21914								
Brevibacterium	lactofermentum				70					
Brevibacterium	lactofermentum				74					
Brevibacterium	lactofermentum				77					
Brevibacterium	lactofermentum	21798								
Brevibacterium	lactofermentum	21799								
Brevibacterium	lactofermentum	21800								
Brevibacterium	lactofermentum	21801								
Brevibacterium	lactofermentum			B11470						
Brevibacterium	lactofermentum			B11471						
Brevibacterium	lactofermentum	21086								
Brevibacterium	lactofermentum	21420								
Brevibacterium	lactofermentum	21086								
Brevibacterium	lactofermentum	31269								
Brevibacterium	linens	9174								
Brevibacterium	linens	19391								
Brevibacterium	linens	8377								
Brevibacterium	paraffinolyticum					11160				
Brevibacterium	spec.						717.73			
Brevibacterium	spec.	14604					717.73			
Brevibacterium	spec.	21860								
Brevibacterium	spec.	21864								
Brevibacterium	spec.	21865								

Corynebacterium	glutamicum	21514																	
Corynebacterium	glutamicum	21516																	
Corynebacterium	glutamicum	21299																	
Corynebacterium	glutamicum	21300																	
Corynebacterium	glutamicum	39684																	
Corynebacterium	glutamicum	21488																	
Corynebacterium	glutamicum	21649																	
Corynebacterium	glutamicum	21650																	
Corynebacterium	glutamicum	19223																	
Corynebacterium	glutamicum	13869																	
Corynebacterium	glutamicum	21157																	
Corynebacterium	glutamicum	21158																	
Corynebacterium	glutamicum	21159																	
Corynebacterium	glutamicum	21355																	
Corynebacterium	glutamicum	31808																	
Corynebacterium	glutamicum	21674																	
Corynebacterium	glutamicum	21562																	
Corynebacterium	glutamicum	21563																	
Corynebacterium	glutamicum	21564																	
Corynebacterium	glutamicum	21565																	
Corynebacterium	glutamicum	21566																	
Corynebacterium	glutamicum	21567																	
Corynebacterium	glutamicum	21568																	
Corynebacterium	glutamicum	21569																	
Corynebacterium	glutamicum	21570																	
Corynebacterium	glutamicum	21571																	
Corynebacterium	glutamicum	21572																	
Corynebacterium	glutamicum	21573																	
Corynebacterium	glutamicum	21579																	
Corynebacterium	glutamicum	19049																	
Corynebacterium	glutamicum	19050																	

Corynebacterium	glutamicum	19051								
Corynebacterium	glutamicum	19052								
Corynebacterium	glutamicum	19053								
Corynebacterium	glutamicum	19054								
Corynebacterium	glutamicum	19055								
Corynebacterium	glutamicum	19056								
Corynebacterium	glutamicum	19057								
Corynebacterium	glutamicum	19058								
Corynebacterium	glutamicum	19059								
Corynebacterium	glutamicum	19060								
Corynebacterium	glutamicum	19185								
Corynebacterium	glutamicum	13286								
Corynebacterium	glutamicum	21515								
Corynebacterium	glutamicum	21527								
Corynebacterium	glutamicum	21544								
Corynebacterium	glutamicum	21492								
Corynebacterium	glutamicum				B8183					
Corynebacterium	glutamicum				B8182					
Corynebacterium	glutamicum				B12416					
Corynebacterium	glutamicum				B12417					
Corynebacterium	glutamicum				B12418					
Corynebacterium	glutamicum				B11476					
Corynebacterium	glutamicum	21608								
Corynebacterium	lilium				P973					
Corynebacterium	nitrilophilus	21419						11594		
Corynebacterium	spec.				P4445					
Corynebacterium	spec.				P4446					
Corynebacterium	spec.	31088								
Corynebacterium	spec.	31089								
Corynebacterium	spec.	31090								
Corynebacterium	spec.	31090								

Corynebacterium	spec.	31090								
Corynebacterium	spec.	15954							20145	
Corynebacterium	spec.	21857								
Corynebacterium	spec.	21862								
Corynebacterium	spec.	21863								

ATCC: American Type Culture Collection, Rockville, MD, USA

FERM: Fermentation Research Institute, Chiba, Japan

NRRL: ARS Culture Collection, Northern Regional Research Laboratory, Peoria, IL, USA

CECT: Coleccion Espanola de Cultivos Tipo, Valencia, Spain

NCIMB: National Collection of Industrial and Marine Bacteria Ltd., Aberdeen, UK

CBS: Centraalbureau voor Schimmelcultures, Baam, NL

NCTC: National Collection of Type Cultures, London, UK

DSMZ: Deutsche Sammlung von Mikroorganismen und Zellkulturen, Braunschweig, Germany

For reference see Sugawara, H. et al. (1993) World directory of collections of cultures of microorganisms: Bacteria, fungi and yeasts (4th edn), World federation for culture collections world data center on microorganisms, Saimata, Japan.



00/02/90

80209/60
OLD S.N 85821

Attorney Docket No.: BGI-124CP

TABLE 4: ALIGNMENT RESULTS

ID #	length	Genbank Hit	Length	Accession	Name of Genbank Hit	Source of Genbank Hit	% homology (GAP)	Date of Deposit
ra00062	1521	GB_HTG2:AC007366	185001	AC007366	Homo sapiens clone NH0501622. *** SEQUENCING IN PROGRESS ***. 3 unordered pieces.	Homo sapiens	39.080	5-Jun-99
ra00084	948	GB_PR3:HSU80741 GB_PL1:BNDNATRNA GB_PR3:HSU80741 GB_GSS9:AC0163721	912 1732 912 388	U80741 X89901 U80741 AC0163721	Homo sapiens CAGH44 mRNA, partial cds. B. nigra DNA for rRNA like gene. Homo sapiens CAGH44 mRNA, partial cds. HS_2245_A1_F07_MF CIT Approved Human Genomic Sperm Library D Homo sapiens genomic clone Plate=2245 Col=13 Row=K, genomic survey sequence.	Homo sapiens Brassica nigra Homo sapiens Homo sapiens	39.264 36.725 38.957 45.066	18-DEC-1997 6-Feb-97 18-DEC-1997 16-OCT-1998
ra00109	735	GB_HTG4:AC007054	171979	AC007054	Drosophila melanogaster chromosome 2 clone BACR45018 (D527) RPCI-98 45.O.18 map 41E-41E strain Y, cn bw sp, *** SEQUENCING IN PROGRESS***. 13 unordered pieces.	Drosophila melanogaster	36.589	13-OCT-1999
		GB_HTG4:AC007054	171979	AC007054	Drosophila melanogaster chromosome 2 clone BACR45018 (D527) RPCI-98 45.O.18 map 41E-41E strain Y, cn bw sp, *** SEQUENCING IN PROGRESS***. 13 unordered pieces.	Drosophila melanogaster	36.589	13-OCT-1999
ra00215	1449	GB_BA1:SC9C7 GB_BA1:SCE94 GB_BA2:AF110185	31360 38532 20302	AL035161 AL049628 AF110185	Streptomyces coelicolor cosmid 9C7. Streptomyces coelicolor cosmid E94. Burkholderia pseudomallei strain 1026b DbhB (dbhB), general secretory pathway protein D (gspD), general secretory pathway protein E (gspE), general secretory pathway protein F (gspF), GspC (gspC), general secretory pathway protein G (gspG), general secretory pathway protein H (gspH), general secretory pathway protein I (gspI), general secretory pathway protein J (gspJ), general secretory pathway protein K (gspK), general secretory pathway protein L (gspL), general secretory pathway protein M (gspM), and general secretory pathway protein N (gspN) genes, complete cds; and unknown genes.	Streptomyces coelicolor Streptomyces coelicolor Burkholderia pseudomallei	44.444 36.313 44.159	12-Jan-99 12-Apr-99 2-Aug-99
ra00289	1299	GB_EST6:N80167	384	N80167	z66g02.s1 Soares fetal liver spleen 1NFLS Homo sapiens cDNA clone IMAGE:297458 Homo sapiens 3', mRNA sequence.	Homo sapiens	40.420	29-MAR-1996
		GB_ST5:G37084 GB_ST5:G37084 GB_BA1:MTCY22D7	384 384 31859	G37084 G37084 Z83866	SHGC-56832 Human Homo sapiens STS genomic, sequence tagged site. SHGC-56832 Human Homo sapiens STS genomic, sequence tagged site. Mycobacterium tuberculosis H37Rv complete genome, segment 133/162.	Homo sapiens Homo sapiens Mycobacterium tuberculosis	40.420 40.420 60.271	30-MAR-1998 30-MAR-1998 17-Jun-98
ra00404	2439	GB_BA1:ECU82598 GB_BA2:AE000165 GB_BA1:SCF43A	136742 12003 35437	U82598 AE000165 AL096837	Escherichia coli genomic sequence of minutes 9 to 12. Escherichia coli K-12 MG1655 section 55 of 400 of the complete genome. Streptomyces coelicolor cosmid F43A.	Escherichia coli Escherichia coli Streptomyces coelicolor	54.256 54.256 36.245	15-Jan-97 12-Nov-98 13-Jul-99
ra00479	2313	GB_GSS2:CNS015U4	1036	AL105910	Drosophila melanogaster genome survey sequence SP6 end of BAC BACN14G08 of DrosBAC library from Drosophila melanogaster (fruit fly), genomic survey sequence.	Drosophila melanogaster	37.573	26-Jul-99
ra00497	420	GB_PR3:HS494O16 GB_BA1:MTCY78	50502 33818	AL117328 Z77165	Human DNA sequence from clone 494O16 on chromosome 22, complete sequence. Mycobacterium tuberculosis H37Rv complete genome, segment 145/162.	Homo sapiens Mycobacterium tuberculosis	36.475 40.250	23-Nov-99 17-Jun-98
		GB_BA2:AF079544	817	AF079544	Mycobacterium avium GroESL operon, partial sequence.	Mycobacterium avium	64.439	16-Aug-98

TABLE 4: ALIGNMENT RESULTS

	GB_BA1:MTGROEOP	2987	X60350	M.tuberculosis groE gene for KCS and 10-kDa products.	Mycobacterium tuberculosis 62,857	23-Apr-92
ra00575						
ra00599 510	GB_GSS10:AQ199703	439	AQ199703	RPC11-46013.TJ RPC1-11 Homo sapiens genomic clone RPC1-11-46013, genomic survey sequence.	Homo sapiens	42,657
	GB_PRR2:AC002127	144165	AC002127	Human BAC clone RG305H12 from 7q21, complete sequence.	Homo sapiens	37,052
	GB_STS:G51234	439	G51234	SHGC-80708 Human Homo sapiens STS genomic, sequence tagged site.	Homo sapiens	42,657
ra00600 1221	GB_BA1:MTCY441	35187	Z80225	Mycobacterium tuberculosis H37Rv complete genome, segment 118/162.	Mycobacterium tuberculosis 56,183	18-Jun-98
	GB_BA1:MSGY223	42061	AD000019	Mycobacterium tuberculosis sequence from clone Y223.	Mycobacterium tuberculosis 37,217	10-DEC-1996
ra00605 1603	GB_BA1:BSUB0014	213420	Z99117	Bacillus subtilis complete genome (section 14 of 21); from 2599451 to 2812870.	Bacillus subtilis	36,553
	GB_BA2:AF069070	2776	AF069070	Endosymbiont of Onchocerca volvulus catalase gene, complete cds.	endosymbiont of Onchocerca volvulus	55,396
	GB_BA1:OVCA1	1845	X82176	Onchocerca volvulus endobacterial mRNA for catalase.	endosymbiont of Onchocerca volvulus	55,396
ra00648 1533	GB_BA1:SC2G5	38404	AL035478	Streptomyces coelicolor cosmid 2G5.	Streptomyces coelicolor	39,530
	GB_HTG1:HS74016	169401	AL110119	Homo sapiens chromosome 21 clone RPC1P70401674 map 21q21, *** SEQUENCING IN PROGRESS ***, in unordered pieces.	Homo sapiens	36,327
	GB_HTG1:HS74016	169401	AL110119	Homo sapiens chromosome 21 clone RPC1P70401674 map 21q21, *** SEQUENCING IN PROGRESS ***, in unordered pieces.	Homo sapiens	36,327
	GB_HTG1:HS74016	169401	AL110119	Homo sapiens chromosome 21 clone RPC1P70401674 map 21q21, *** SEQUENCING IN PROGRESS ***, in unordered pieces.	Homo sapiens	36,327
ra00764 1239	GB_EST36:AI898007	609	AI898007	EST267450 tomato ovary, TAMU Lycopersicon esculentum cDNA clone cLED31K22, mRNA sequence.	Lycopersicon esculentum	34,323
	GB_BA2:PAU93274	8008	U93274	Pseudomonas aeruginosa YalE (yalE), LeuB (leuB), Asd (asd), FimV (fimV), and HisT (hisT) genes, complete cds; TrpF (trpF) gene, partial cds; and unknown gene.	Pseudomonas aeruginosa	35,895
	GB_BA2:PAU93274	8008	U93274	Pseudomonas aeruginosa YalE (yalE), LeuB (leuB), Asd (asd), FimV (fimV), and HisT (hisT) genes, complete cds; TrpF (trpF) gene, partial cds; and unknown gene.	Pseudomonas aeruginosa	41,417
ra00803 1353	GB_IN2:CELH34C03	27748	AF100662	Caenorhabditis elegans cosmid H34C03.	Caenorhabditis elegans	34,152
	GB_HTG2:AC007905	100722	AC007905	Homo sapiens chromosome 16q24.3 clone PAC 754F23, *** SEQUENCING IN PROGRESS ***, 33 unordered pieces.	Homo sapiens	37,472
	GB_HTG2:AC007905	100722	AC007905	Homo sapiens chromosome 16q24.3 clone PAC 754F23, *** SEQUENCING IN PROGRESS ***, 33 unordered pieces.	Homo sapiens	37,472
ra00810 324	GB_BA1:MTV15C10	33050	Z95436	Mycobacterium tuberculosis H37Rv complete genome, segment 154/162.	Mycobacterium tuberculosis 34,615	17-Jun-98
	GB_BA1:MLCB2548	38916	AL023093	Mycobacterium leprae cosmid B2548.	Mycobacterium leprae	34,615
	GB_BA1:ECOUW76	225419	U00039	E. coli chromosomal region from 76.0 to 81.5 minutes.	Escherichia coli	52,997
ra00829 2463	GB_BA1:SC5C7	41906	AL015115	Streptomyces coelicolor cosmid 5C7.	Streptomyces coelicolor	65,269
	GB_BA1:SC5F2A	40105	AL049587	Streptomyces coelicolor cosmid 5F2A.	Streptomyces coelicolor	37,490
	GB_BA1:STMDRRC	3374	L76359	Streptomyces peucetius daunorubicin resistance protein (drc) gene, complete cds.	Streptomyces peucetius	55,279
ra00843 468	GB_BA1:MTCY9C4	15916	Z77250	Mycobacterium tuberculosis H37Rv complete genome, segment 113/162.	Mycobacterium tuberculosis 40,000	17-Jun-98

TABLE 4: ALIGNMENT RESULTS

	GB_BA1:MTCY9C4	15916	Z77250	Mycobacterium tuberculosis H37Rv complete genome; segment 113/162.	Mycobacterium tuberculosis 37.773	17-Jun-98
ra00858 568	GB_BA1:SCC54	30753	AL035591	Streptomyces coelicolor cosmid C54.	Streptomyces coelicolor 39.602	11-Jun-99
	GB_EST18:N96610	547	N96610	21285 Lambda-PRL1 Arabidopsis thaliana cDNA clone F10G3T7, mRNA sequence.	Arabidopsis thaliana 37.801	5-Jan-98
	GB_EST18:T45493	436	T45493	8756 Lambda-PRL2 Arabidopsis thaliana cDNA clone 133C14T7, mRNA sequence.	Arabidopsis thaliana 34.194	4-Aug-98
ra00886 1269	GB_BA1:SYCSLLH	132106	DC4006	Synechocystis sp. PCC6803 complete genome, 25/27, 3138604-3270709.	Synechocystis sp. 37.459	13-Feb-99
	GB_BA1:SCDNAJ	5611	X77458	S.coelicolor dnaK, gprE and dnaJ genes.	Streptomyces coelicolor 49.744	21-Nov-96
	GB_BA1:STMDNAK	4648	L46700	Streptomyces coelicolor (strain A3(2)) dnaK operon encoding molecular chaperones (dnaK, dnaJ, gprE and hspR genes, complete cds).	Streptomyces coelicolor 49.583	22-Nov-96
ra00900 975	GB_BA2:ECOWW67_0	110000	U18997	Escherichia coli K-12 chromosomal region from 67.4 to 76.0 minutes.	Escherichia coli 38.314	U18997
	GB_BA2:ECOWW67_0	110000	U18997	Escherichia coli K-12 chromosomal region from 67.4 to 76.0 minutes.	Escherichia coli 37.759	U18997
	GB_BA2:AE000393	10516	AE000393	Escherichia coli K-12 MG1655 section 283 of 400 of the complete genome.	Escherichia coli 38.314	12-Nov-98
ra00901 537	GB_HTG3:AC010757	175571	AC010757	Homo sapiens chromosome 18 clone 128_C_18 map 18, *** SEQUENCING IN PROGRESS *** , 20 unordered pieces.	Homo sapiens 34.857	22-Sep-99
	GB_HTG3:AC010757	175571	AC010757	Homo sapiens chromosome 18 clone 128_C_18 map 18, *** SEQUENCING IN PROGRESS *** , 20 unordered pieces.	Homo sapiens 34.857	22-Sep-99
	GB_HTG3:AC011283	87295	AC011283	Homo sapiens clone MS2016A09, *** SEQUENCING IN PROGRESS *** , 1 unordered pieces.	Homo sapiens 35.448	07-OCT-1999
ra00981 753	GB_OV:GGA245664	512	AJ245664	Gallus gallus partial mRNA for ATP-citrate lyase (ACL gene).	Gallus gallus 37.538	28-Sep-99
	GB_PL2:AC007887	159434	AC007887	Genomic sequence for Arabidopsis thaliana BAC F1504 from chromosome I, complete sequence.	Arabidopsis thaliana 37.600	04-OCT-1999
	GB_GSS1:CNS00RNV 542		AL087338	Arabidopsis thaliana genome survey sequence T7 end of BAC F14D7 of IGF library from Arabidopsis thaliana strain Columbia of Arabidopsis thaliana, genomic survey sequence.		28-Jun-99
ra00995 864	GB_EST29:AI553951	450	AI553951	te54401.x1 Soares, NFL_T_GBC_S1 Homo sapiens cDNA clone IMAGE:2090497.3 similar to gb:X02067 H.sapiens mRNA for 7SL RNA pseudogene (HUMAN)., mRNA sequence.	Homo sapiens 42.627	13-Apr-99
	GB_PR3:AC003029	139166	AC003029	Homo sapiens Chromosome 12q24 PAC RPC13-462E2 (Roswell Park Cancer Institute Human PAC library) complete sequence.	Homo sapiens 38.915	17-Sep-98
	GB_BA1:EAY14603	4479	Y14603	Erwinia amylovora srfA, srfE, srfB, srfD, srfM and srfR genes.	Erwinia amylovora 37.694	6-Jan-98
ra00996 864	GB_BA2:AE001001	10730	AE001001	Archaeoglobus fulgidus section 106 of 172 of the complete genome.	Archaeoglobus fulgidus 41.078	15-DEC-1997
	GB_EST30:AV018764	242	AV018764	AV018764 Mus musculus 18-day embryo C57BL/6J Mus musculus cDNA clone 1190006M16, mRNA sequence.	Mus musculus 39.669	28-Aug-99
	GB_GSS3:B24189	377	B24189	F19E161F IGF Arabidopsis thaliana genomic clone F19E16, genomic survey sequence. Arabidopsis thaliana		10-OCT-1997
ra01010 1242	GB_OV:AF007068	356	AF007068	Columnix columnix any/alkylamine N-acyetyltransferase mRNA, partial cds.	Columnix columnix 46.629	12-Jul-97
	GB_EST10:AA166324	514	AA166324	ms50c09.r1 Lite Tech mouse embryo 13 5dpc 10666014 Mus musculus cDNA clone IMAGE:614992.5 similar to SW:NEST_RAT P21263 NESTIN. ., mRNA sequence.	Mus musculus 38.677	19-DEC-1996
	GB_EST7:W89968	46	W89968	mt64g11.r1 Soares mouse embryo NbMc13.5 14.5 Mus musculus cDNA clone IMAGE:419108.5 similar to SW:NEST_RAT P21263 NESTIN.[1]., mRNA sequence.	Mus musculus 58.696	12-Sep-96
ra01051 732	GB_GSS12:AQ381423	579	AQ381423	RPC11-1-135F10.TJ RPC1-11 Homo sapiens genomic clone RPC1-11-135F10, genomic survey sequence.	Homo sapiens 37.651	21-MAY-1999
	GB_HTG6:AC010901	206121	AC010901	Homo sapiens clone RP11-544J22, WORKING DRAFT SEQUENCE, 1 unordered pieces.	Homo sapiens 36.011	04-DEC-1999

TABLE 4: ALIGNMENT RESULTS

ra01052	432	GB_IN1:CELC13D9	43487	AF016420	Caenorhabditis elegans cosmid C13D9.	Caenorhabditis elegans	39,344	2-Aug-97
		GB_IN1:CELC13D9	43487	AF016420	Caenorhabditis elegans cosmid C13D9.	Caenorhabditis elegans	38,780	2-Aug-97
ra01053	543	GB_OV:CHKMAFG1	1316	D28601	Chicken novel mat-related gene mafG encoding bZip nuclear protein MafG, promoter region and exon 1.	Gallus gallus	39,205	7-Feb-99
		GB_HTG6:AC010765	146468	AC010765	Homo sapiens clone RP11-115N6, *** SEQUENCING IN PROGRESS ***; 26 unordered Homo sapiens pieces.	Homo sapiens	32,961	07-DEC-1999
		GB_HTG6:AC010765	146468	AC010765	Homo sapiens clone RP11-115N6, *** SEQUENCING IN PROGRESS ***; 26 unordered Homo sapiens pieces.	Homo sapiens	38,476	07-DEC-1999
ra01054	612	GB_PL1:PHNPNGLP	962	DA5425	Pharbitis nil mRNA for Pharbitis nil Germin-like protein precursor, complete cds.	Ipomoea nil	42,925	10-Feb-99
		GB_HTG2:HSJ402N21	170302	AL049553	Homo sapiens chromosome 6 clone RP3.402N21 map p21.1-21.31, ***SEQUENCING IN PROGRESS ***; in unordered pieces.	Homo sapiens	36,825	03-DEC-1999
		GB_HTG2:HSJ402N21	170302	AL049553	Homo sapiens chromosome 6 clone RP3.402N21 map p21.1-21.31, ***SEQUENCING IN PROGRESS ***; in unordered pieces.	Homo sapiens	36,825	03-DEC-1999
ra01217	723	GB_IN2:CELF18A12	29784	AF016688	Caenorhabditis elegans cosmid F18A12.	Caenorhabditis elegans	35,794	08-OCT-1999
		GB_IN2:CELF18A12	29784	AF016688	Caenorhabditis elegans cosmid F18A12.	Caenorhabditis elegans	40,625	08-OCT-1999
		GB_RO:MUSMCFTR	6304	M60493	Mouse cystic fibrosis transmembrane conductance regulator (CFTR) mRNA, complete cds.	Mus musculus	37,793	10-Jun-94
ra01320	1770	GB_BA2:AF031037	1472	AF031037	Neisseria meningitidis chloramphenicol acetyltransferase gene, complete cds.	Neisseria meningitidis	35,014	21-Apr-98
		GB_HTG1:PFMAL13PA 80518	AL109815	AL109815	Plasmidium falciplarum chromosome 13 strain 3D7, *** SEQUENCING IN PROGRESS ***; in unordered pieces.	Plasmidium falciplarum	17,697	19-Aug-99
		GB_HTG1:PFMAL13PA 80518	AL109815	AL109815	Plasmidium falciplarum chromosome 13 strain 3D7, *** SEQUENCING IN PROGRESS ***; in unordered pieces.	Plasmidium falciplarum	17,697	19-Aug-99
ra01345	1575	GB_PR3:AC005224	166687	AC005224	Homo sapiens chromosome 17, clone hRPK.214_O_1, complete sequence.	Homo sapiens	38,195	14-Aug-98
		GB_PR3:AC005224	166687	AC005224	Homo sapiens chromosome 17, clone hRPK.214_O_1, complete sequence.	Homo sapiens	36,611	14-Aug-98
		GB_HTG3:AC011500_1300851	AC011500	AC011500	Homo sapiens chromosome 19 clone CIT9/78SKB_60E11, *** SEQUENCING IN PROGRESS ***; 246 unordered pieces.	Homo sapiens	36,446	AC011500
ra01407	1014	GB_HTG3:AC010831	70233	AC010831	Homo sapiens clone 6_L_24, LOW-PASS SEQUENCE SAMPLING.	Homo sapiens	35,764	23-Sep-99
		GB_HTG3:AC010831	70233	AC010831	Homo sapiens clone 6_L_24, LOW-PASS SEQUENCE SAMPLING.	Homo sapiens	35,764	23-Sep-99
		GB_PR3:AC004058	38400	AC004058	Homo sapiens chromosome 4 clone B241P19 map 4q25, complete sequence.	Homo sapiens	40,778	30-Sep-98
ra01408	324	GB_PR4:AF152365	246546	AF152365	Homo sapiens constitutive fragile region FRA3B sequence.	Homo sapiens	41,234	1-Aug-99
		GB_HTG3:AC007890	121256	AC007890	Drosophila melanogaster chromosome 3 clone BACR02G21 (D722) RPCL:98.02.G.21 map 90E-91A strain y, cn bw sp, *** SEQUENCING IN PROGRESS***; 89 unordered pieces.	Drosophila melanogaster	39,432	3-Sep-99
		GB_HTG3:AC007890	121256	AC007890	Drosophila melanogaster chromosome 3 clone BACR02G21 (D722) RPCL:98.02.G.21 map 90E-91A strain y, cn bw sp, *** SEQUENCING IN PROGRESS***; 89 unordered pieces.	Drosophila melanogaster	39,432	3-Sep-99
ra01524	1566	GB_BA1:BSUB0015	218410	Z99118	Bacillus subtilis complete genome (section 15 of 21): from 2795131 to 3013540.	Bacillus subtilis	38,201	26-Nov-97
		GB_HTG2:AC008260	107439	AC008260	Drosophila melanogaster chromosome 2 clone BACR13J10 (D924) RPCL:98.13.J.10 map 47B-47C strain y, cn bw sp, *** SEQUENCING IN PROGRESS***; 82 unordered pieces.	Drosophila melanogaster	38,302	2-Aug-99

TABLE 4: ALIGNMENT RESULTS

GB_HTG2:AC008260	107439	AC008260	Drosophila melanogaster chromosome 2 clone BACR13J10 (D924) RPC1-98 13.J.10 map 47B-47C strain y; cn bw sp. *** SEQUENCING IN PROGRESS ***. 82 unordered pieces.	Drosophila melanogaster	38,302	2-Aug-99
GB_PR4:AF111170	148083	AF111170	Homo sapiens 14q32 Jagged2 gene, complete cds; and unknown gene.	Homo sapiens	37,873	14-Jul-99
GB_PR4:AF111170	148083	AF111170	Homo sapiens 14q32 Jagged2 gene, complete cds; and unknown gene.	Homo sapiens	40,220	14-Jul-99
GB_BA1:AEY13732	6740	Y13732	Alcaligenes eutrophus genes for ureases, ureD1, ureD2, ureA, ureB, and ORF1, ORF2.	Ralstonia eutropha	42,960	23-Sep-97
GB_BA2:AF088857	2908	AF088857	Vogesella indigofera indigoidine biosynthesis regulatory locus, complete sequence.	Vogesella indigofera	37,626	10-Sep-99
GB_IN1:CEM04D8	21552	Z32682	Caenorhabditis elegans cosmid M04D8, complete sequence.	Caenorhabditis elegans	37,237	23-Nov-98
GB_EST25:AI281910	276	AI281910	q182d04.x1 NC1_CGAP_Co14 Homo sapiens cDNA clone IMAGE:1961767 3'. mRNA sequence.	Homo sapiens	38,406	21-DEC-1998
GB_BA1:CGU43535	2531	UK3535	Corynebacterium glutamicum multidrug resistance protein (cmr) gene, complete cds.	Corynebacterium glutamicum	99,933	9-Apr-97
GB_HTG3:AC009213	114735	AC009213	Drosophila melanogaster chromosome 3 clone BACR09F18 (D812) RPC1-98 09.F.18 map 98D-98D strain y; cn bw sp. *** SEQUENCING IN PROGRESS ***. 109 unordered pieces.	Drosophila melanogaster	36,111	23-Aug-99
GB_HTG3:AC009213	114735	AC009213	Drosophila melanogaster chromosome 3 clone BACR09F18 (D812) RPC1-98 09.F.18 map 98D-98D strain y; cn bw sp. *** SEQUENCING IN PROGRESS ***. 109 unordered pieces.	Drosophila melanogaster	36,111	23-Aug-99
GB_PL1:AB017159	1859	AB017159	Daucus carota mRNA for citrate synthase, complete cds.	Daucus carota	39,537	01-MAY-1999
GB_PR1:HUNGNO548	23142	D26607	Homo sapiens endothelial nitric oxide synthase gene, complete cds.	Homo sapiens	36,419	13-Jul-99
GB_HTG3:AC011234	154754	AC011234	Homo sapiens clone NH0166D23, *** SEQUENCING IN PROGRESS ***. 7 unordered pieces.	Homo sapiens	36,317	04-OCT-1999
GB_HTG3:AC009450	124337	AC009450	Homo sapiens chromosome 9 clone 30_C_23 map 9, *** SEQUENCING IN PROGRESS ***. 20 unordered pieces.	Homo sapiens	35,303	22-Aug-99
GB_HTG3:AC009450	124337	AC009450	Homo sapiens chromosome 9 clone 30_C_23 map 9, *** SEQUENCING IN PROGRESS ***. 20 unordered pieces.	Homo sapiens	35,303	22-Aug-99
GB_HTG3:AC009919	134724	AC009919	Homo sapiens clone 115_L_23, LOW-PASS SEQUENCE SAMPLING.	Homo sapiens	35,409	8-Sep-99
GB_BA1:ECONELUC	1676	M84026	E. coli protein p7 (neu C) gene, complete cds.	Escherichia coli	35,189	26-Apr-93
GB_HTG2:AC007853	116280	AC007853	Drosophila melanogaster chromosome 3 clone BACR03L02 (D766) RPC1-98 03.L.2 map 96B-96C strain y; cn bw sp. *** SEQUENCING IN PROGRESS ***. 80 unordered pieces.	Drosophila melanogaster	34,365	2-Aug-99
GB_HTG2:AC007853	116280	AC007853	Drosophila melanogaster chromosome 3 clone BACR03L02 (D766) RPC1-98 03.L.2 map 96B-96C strain y; cn bw sp. *** SEQUENCING IN PROGRESS ***. 80 unordered pieces.	Drosophila melanogaster	34,365	2-Aug-99
GB_HTG4:AC010037	166249	AC010037	Drosophila melanogaster chromosome 3L/66B6 clone RPC198-6E4, *** SEQUENCING IN PROGRESS ***. 52 unordered pieces.	Drosophila melanogaster	38,534	16-OCT-1999
GB_HTG4:AC010037	166249	AC010037	Drosophila melanogaster chromosome 3L/66B6 clone RPC198-6E4, *** SEQUENCING IN PROGRESS ***. 52 unordered pieces.	Drosophila melanogaster	38,534	16-OCT-1999
GB_PR4:AC005552	167228	AC005552	Homo sapiens chromosome 17, clone hRPK_212_E_8, complete sequence.	Homo sapiens	36,249	26-Nov-98
GB_PR1:HS169C8F	245	Z57239	H. sapiens CPG island DNA genomic MseI fragment, clone 169c8, forward read cpg169c8.t1a.	Homo sapiens	45,679	18-OCT-1995
GB_BA1:SERATTBXIS	3255	L11597	Saccharopolyspora erythraea excisionase (xis) gene, integrase (int) gene, complete cds's and attB site.	Saccharopolyspora erythraea	36,232	6-Jul-94

TABLE 4: ALIGNMENT RESULTS

ra02060	GB_EST7:W97557	267	W97557	mt8a09.r1 Soares mouse embryo NbME13.5 14.5 Mus musculus cDNA clone IMAGE:422296 5', mRNA sequence.	Mus musculus	42,969	16-Jul-96
ra02087 1470	GB_PR3:AC005544	169045	AC005544	Homo sapiens chromosome 17, clone hRPK_349_A_8, complete sequence.	Homo sapiens	35,724	25-Sep-98
	GB_PL1:ATF20B18	104738	AL049483	Arabidopsis thaliana DNA chromosome 4, BAC clone F20B18 (ESSA project).	Arabidopsis thaliana	35,890	24-MAR-1999
	GB_PL2:ATT29K17	89904	AL049171	Arabidopsis thaliana DNA chromosome 4, BAC clone (ESSA project).	Arabidopsis thaliana	38,128	27-Aug-99
ra02088 1338	GB_HTG3:AC008697	167932	AC008697	Homo sapiens chromosome 5 clone CIT978SKB_70D3, *** SEQUENCING IN PROGRESS ***. 54 unordered pieces.	Homo sapiens	36,662	3-Aug-99
	GB_HTG3:AC008697	167932	AC008697	Homo sapiens chromosome 5 clone CIT978SKB_70D3, *** SEQUENCING IN PROGRESS ***. 54 unordered pieces.	Homo sapiens	36,662	3-Aug-99
	GB_HTG3:AC008703	213971	AC008703	Homo sapiens chromosome 5 clone CIT978SKB_76P12, *** SEQUENCING IN PROGRESS ***. 54 unordered pieces.	Homo sapiens	34,768	3-Aug-99
ra02159 636	GB_BA2:AF049897	9196	AF049897	Corynebacterium glutamicum N-acetylglutamylphosphate reductase (argC), ornithine acetyltransferase (argJ), N-acetylglutamate kinase (argB), acetylornithine transaminase (argD), ornithine carbamoyltransferase (argF), arginine repressor (argR), argininosuccinate synthase (argG), and argininosuccinate lyase (argH) genes, complete cds.	Corynebacterium glutamicum	99,843	1-Jul-98
	GB_BA2:AF031518	2045	AF031518	Corynebacterium glutamicum ornithine carbamoyltransferase (argF) gene, complete cds.	Corynebacterium glutamicum	88,679	5-Jan-99
	GB_BA2:AF041436	516	AF041436	Corynebacterium glutamicum arginine repressor (argR) gene, complete cds.	Corynebacterium glutamicum	100,000	5-Jan-99
ra02184 504	GB_BA1:BSZ92953	8164	Z92953	B subtilis yws[A,B,C] genes and rbs[A,C,D,K,R] genes.	glutamicum	38,951	24-Jun-98
	GB_EST36:A1878071	593	A1878071	tc57a12.y1 Zebrafish Washu MPMWG EST Danio rerio cDNA 5' similar to TR:Q13151 Q13151 HETEROGENEOUS NUCLEAR RIBONUCLEOPROTEIN A0 ;, mRNA sequence.	Bacillus subtilis	36,774	21-Jul-99
	GB_EST37:A1958166	641	A1958166	tc91f01.y1 Zebrafish Washu MPMWG EST Danio rerio cDNA 5' similar to TR:Q13151 Q13151 HETEROGENEOUS NUCLEAR RIBONUCLEOPROTEIN A0 ;, mRNA sequence.	Danio rerio	36,774	20-Aug-99
ra02200 1233	GB_PR3:HS4494O16	50502	AL117328	Human DNA sequence from clone 494O16 on chromosome 22, complete sequence.	Homo sapiens	38,648	23-Nov-99
	GB_HTG2:AC008161	158440	AC008161	Mus musculus clone 182_H_5, *** SEQUENCING IN PROGRESS ***. 29 unordered pieces.	Mus musculus	35,938	28-Jul-99
	GB_HTG2:AC008161	158440	AC008161	Mus musculus clone 182_H_5, *** SEQUENCING IN PROGRESS ***. 29 unordered pieces.	Mus musculus	35,938	28-Jul-99
ra02201 486	GB_EST4:H16949	465	H16949	ym34a11.r1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:50010 5', mRNA sequence.	Homo sapiens	38,267	29-Jun-95
	GB_EST4:H16949	465	H16949	ym34a11.r1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:50010 5', mRNA sequence.	Homo sapiens	36,552	29-Jun-95
ra02202 762	GB_IN1:CELCA1A3	37149	U41541	Caenorhabditis elegans cosmid C41A3.	Caenorhabditis elegans	41,678	08-DEC-1995
	GB_EST33:AV080151	236	AV080151	AV080151 Mus musculus stomach C57BL/6J adult Mus musculus cDNA clone 2210413B04, mRNA sequence.	Mus musculus	43,348	25-Jun-99
	GB_GSS5:AQ766877	545	AQ766877	HS_2017_B2_B08_MR CIT Approved Human Genomic Sperm Library D Homo sapiens genomic clone Plate=2017 Col=16 Row=D, genomic survey sequence.	Homo sapiens	35,568	28-Jul-99

TABLE 4: ALIGNMENT RESULTS

ra02205	1002	GB_HTG2:AC005959	127587	AC005959	Homo sapiens *** SEQUENCING IN PROGRESS *** 2 ordered pieces.	Homo sapiens	40,310	11-Nov-98
		GB_HTG2:AC005959	127587	AC005959	Homo sapiens, *** SEQUENCING IN PROGRESS *** 2 ordered pieces.	Homo sapiens	40,310	11-Nov-98
		GB_IN1:BRPTUBBA	4571	M36380	B.pahangi beta-tubulin gene, complete cds.	Brugia pahangi	37,703	26-Apr-93
ra02305	975	GB_RO:MUSPARR	1140	D50872	Mouse gene for platelet activating factor receptor, complete cds.	Mus musculus	38,420	10-Feb-99
		GB_PR3:HUMARL1A	1008	L28997	Homo sapiens ARL1 mRNA, complete cds.	Homo sapiens	42,188	13-Jan-95
		GB_BA1:MLCB2533	40245	AL035310	Mycobacterium leprae cosmid B2533.	Mycobacterium leprae	42,000	27-Aug-99
ra02431	899	GB_EST4:H35255	407	H35255	EST111890 Rat PC-12 cells, NGF-treated (9 days) Rattus sp. cDNA clone RPNCO03, mRNA sequence.	Rattus sp.	39,098	2-Apr-98
		GB_HTG1:HS791K14	155318	AL035685	Homo sapiens chromosome 20 clone RP4-791K14, *** SEQUENCING IN PROGRESS *** in unordered pieces.	Homo sapiens	39,456	23-Nov-99
		GB_HTG1:HS791K14	155318	AL035685	Homo sapiens chromosome 20 clone RP4-791K14, *** SEQUENCING IN PROGRESS *** in unordered pieces.	Homo sapiens	39,456	23-Nov-99
ra02446	558	GB_BA2:AF036166	895	AF036166	Xanthomonas campestris organic hydroperoxide resistance protein (ohr) gene, completeXanthomonas campestris cds.	Xanthomonas campestris	49,369	19-MAY-1998
		GB_EST5:N25122	620	N25122	yx19d10.r1 Soares melanocyte 2NbhM Homo sapiens cDNA clone IMAGE:262195 5', mRNA sequence.	Homo sapiens	35,417	28-DEC-1995
		GB_EST5:N25122	620	N25122	yx19d10.r1 Soares melanocyte 2NbhM Homo sapiens cDNA clone IMAGE:262195 5', mRNA sequence.	Homo sapiens	37,172	28-DEC-1995
ra02541	1308	GB_BA2:DPJ93358	1267	U93358	Deinococcus proteolyticus 40 kDa heat shock chaperone protein (dnaJ) gene, complete Deinococcus proteolyticus cds.	Deinococcus proteolyticus	42,115	17-Jan-98
		GB_EST30:A1658096	343	A1658096	fc14c09.y1 Zebratish WashU MPIMG EST Danio rerio cDNA 5' similar to SW:DNJ2_HUMAN P31689 DNAU PROTEIN HOMOLOG 2. ; mRNA sequence.	Danio rerio	52,059	06-MAY-1999
		GB_EST37:A1959242	545	A1959242	fg25h11.y1 Zebratish WashU MPIMG EST Danio rerio cDNA 5' similar to SW:DNJ2_HUMAN P31689 DNAU PROTEIN HOMOLOG 2. ; mRNA sequence.	Danio rerio	45,438	20-Aug-99
ra02542	777	EM_PAT:E10832	1856	E10832	DNA encoding Dnak protein which is one of heat shock protein from	Corynebacterium glutanicum	99,000	08-OCT-1997
		GB_EST24:Z82017	396	Z82017	SSZ82017 Porcine small intestine cDNA library Sus scrofa cDNA clone c12c06 5' similarSus scrofa to eukaryotic initiation factor 4 gamma. mRNA sequence.	Felis catus	39,409	14-OCT-1993
ra02543	1977	GB_OM:CATERYTHRO 681	L10606	L10606	Cat erythropoietin mRNA, 3' end.	Corynebacterium glutanicum	97,306	08-OCT-1997
		EM_PAT:E10832	1856	E10832	DNA encoding Dnak protein which is one of heat shock protein from	Corynebacterium glutanicum	97,306	(Rel. 52, Created)
		GB_BA1:MPHSP70	2179	X59437	M.paratuberculosis gene for 70 kD heat shock protein.	Mycobacterium avium subsp. paratuberculosis	73,404	23-Apr-92
ra02586	393	GB_BA1:MTY13E10	35019	Z85324	Mycobacterium tuberculosis H37Rv complete genome, segment 18/162.	Mycobacterium tuberculosis	72,028	17-Jun-98
		GB_IN2:AC006472	156362	AC006472	Drosophila melanogaster, chromosome 2R, region 45E1-46A2, BAC clone BACR48G21, complete sequence.	Drosophila melanogaster	37,958	30-Jan-99
		GB_HTG4:AC010020	106541	AC010020	Drosophila melanogaster chromosome 3L/66D10 clone RPC198-2613, *** SEQUENCINGDrosophila melanogaster IN PROGRESS *** 55 unordered pieces.	Drosophila melanogaster	37,333	16-OCT-1999
		GB_HTG4:AC010020	106541	AC010020	Drosophila melanogaster chromosome 3L/66D10 clone RPC198-2613, *** SEQUENCINGDrosophila melanogaster IN PROGRESS *** 55 unordered pieces.	Drosophila melanogaster	37,333	16-OCT-1999
ra02587	2214	GB_BA1:MLCL622	42498	Z95398	Mycobacterium leprae cosmid L622.	Mycobacterium leprae	39,848	24-Jun-97
		GB_RO:AF074879	3316	AF074879	Rattus norvegicus testis-specific protein TSPY gene, complete cds.	Rattus norvegicus	35,830	6-Jul-99

TABLE 4: ALIGNMENT RESULTS

rx03217 331	GB_RQ:RNJ001380	2641	AJ001380	Rattus norvegicus Tspy partial genomic sequence, exons 1-6.	Rattus norvegicus	37,702	29-Jun-98
	GB_BA1:MLCB2548	38916	AL023093	Mycobacterium leprae cosmid B2548.	Mycobacterium leprae	37,888	-27-Aug-99
	GB_HTG2:HSJ662M14	174772	AL079336	Homo sapiens chromosome 20 clone RP4-662M14, *** SEQUENCING IN PROGRESS	Homo sapiens	36,420	4-Feb-00
	GB_HTG2:HSJ662M14	174772	AL079336	*** 10 unordered pieces. Homo sapiens chromosome 20 clone RP4-662M14, *** SEQUENCING IN PROGRESS	Homo sapiens	35,962	4-Feb-00